

DESREM™

Lyophilized Powder for Injection for IV Infusion

chemistries, hematology, ALT, AST, bilirubin, and alkaline phosphatase; renal function tests (creatinine and creatinine clearance).

8. DRUG INTERACTIONS

Drug-drug interaction trials of remdesivir and other concomitant medications have not been conducted in humans. In vitro, remdesivir is a substrate for drug metabolizing enzymes CYP2C8, CYP2D6, and CYP3A4, and is a substrate for Organic Anion Transporting Polypeptides 1B1 (OATP1B1) and P-glycoprotein (Pgp) transporters. In vitro, remdesivir is an inhibitor of CYP3A4, OATP1B1, OATP1B3, BSEP, MRP4, and NTCP. The clinical relevance of these in vitro assessments has not been established.

9. USE IN SPECIFIC POPULATIONS

9.1 Pregnancy

Risk Summary

No adequate and well-controlled studies of remdesivir use in pregnant women have been conducted. Remdesivir should be used during pregnancy only if the potential benefit justifies the potential risk for the mother and the fetus. In nonclinical reproductive toxicity studies, remdesivir demonstrated no adverse effect on embryofetal development when administered to pregnant animals at systemic exposures (AUC) of the predominant circulating metabolite of remdesivir (GS-441524) that were 4 times (rats and rabbits) the exposure in humans at the recommended human dose (RHD).

Animal Data

Remdesivir was administered via intravenous injection to pregnant rats and rabbits (up to 20 mg/kg/day) on Gestation Days 6 through 17, and 7 through 20, respectively, and also to rats from Gestation Day 6 to Lactation/Post-partum Day 20. No adverse effects on embryo-fetal (rats and rabbits) or pre/postnatal (rats) development were observed in rats and rabbits at nontoxic doses in pregnant animals. During organogenesis, exposures to the predominant circulating metabolite (GS-441524) were 4 (rats and rabbits) times higher than the exposure in humans at the RHD. In a pre/postnatal development study, exposures to the predominant circulating metabolite of remdesivir (GS-441524) were similar to the human exposures at the RHD.

9.2 Nursing Mothers

Risk Summary

There is no information regarding the presence of remdesivir in human milk, the effects on the breastfed infant, or the effects on milk production. In animal studies, remdesivir and metabolites have been detected in the nursing pups of mothers given remdesivir, likely due to the presence of remdesivir in milk. Because of the potential for viral transmission to SARS-CoV-2-negative infants and adverse reactions from the drug in breastfeeding infants, the developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for remdesivir and any potential adverse effects on the breastfed child from remdesivir or from the underlying maternal condition.

Animal Data

Remdesivir and its metabolites were detected in the plasma of nursing rat pups, likely due to the presence of remdesivir and/or its metabolites in milk, following daily intravenous administration of remdesivir to pregnant mothers from Gestation Day 6 to Lactation Day 20. Exposures in nursing pups were approximately 1% that of maternal exposure on lactation day 10.

9.3 Pediatric Use

The safety and effectiveness of remdesivir for treatment of COVID-19 have not been assessed in pediatric patients. Dosing instructions for pediatric patients were derived based on pharmacokinetic data from adult healthy volunteers and *in vitro* data for remdesivir and other similar compounds, as part of the PBPK modeling and simulation approach which accounts for age-dependent changes in metabolism, distribution, and elimination of remdesivir.

For pediatric patients with body weight between 3.5 kg to <40 kg, use remdesivir for injection, 100 mg, lyophilized powder only [see Dosage and Administration (2.3 and 2.8)].

Pediatric patients (>28 days) must have creatinine clearance determined and fullterm neonates (≥7 days to <28 days) must have serum creatinine determined before dosing. Pediatric patients should be monitored for renal function and consideration given for stopping therapy in the setting of substantial decline. The use of remdesivir is not recommended in pediatric patients (>28 days old) with eGFR <30 mL/min and in full-term neonates (≥7 days and ≤28 days old) with serum creatinine clearance ≥1 mg/dL unless the potential benefit outweighs the potential risk.

Because the excipient sulfolbutylether-β-cyclodextrin sodium salt (SBECD) is renally cleared and accumulates in patients with decreased renal function, administration of drugs formulated with SBECD (such as remdesivir) is not recommended in adults and pediatric patients (>28 days old) with eGFR less than 30 mL per minute or in full-term neonates (≥7 days and ≤28 days old) with serum creatinine clearance ≥1 mg/dL unless the potential benefit outweighs the potential risk.

9.4 Geriatric Use

The pharmacokinetics of remdesivir have not been evaluated in patients >65 years of age. In general, appropriate caution should be exercised in the administration of remdesivir and monitoring of elderly patients, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

9.5 Renal Impairment

Use in patients with renal impairment are based on potential risk and potential benefit considerations. Patients with eGFR greater than or equal to 30 mL/min are reported to have received remdesivir for treatment of COVID-19 with no dose adjustment of remdesivir. All patients must have an eGFR determined before dosing.

9.6 Hepatic Impairment

The pharmacokinetics of remdesivir have not been evaluated in patients with hepatic impairment. It is not known if dosage adjustment is needed in patients with hepatic impairment and remdesivir should only be used in patients with hepatic impairment if the potential benefit outweighs the potential risk [see Warnings and Precautions (5.2)].

Hepatic laboratory testing should be performed in all patients prior to starting remdesivir and daily while receiving remdesivir.


10. OVERDOSAGE

There is no human experience of acute overdosage with remdesivir. Treatment of overdose with remdesivir should consist of general supportive measures including monitoring of vital signs and observation of the clinical status of the patient. There is no specific antidote for overdose with remdesivir.

11. PRODUCT DESCRIPTION

Remdesivir is a nucleoside ribonucleic acid (RNA) polymerase inhibitor.

The chemical name for remdesivir is 2-ethylbutyl *N*-(*S*)-[2-*C*-(4aminopyrrolo[2,1-*f*][1,2,4]triazin-7-yl)-2,5-anhydro-d-altronitril-6-*Oy*]phenoxyphosphoryl]-L-alaninate. It has a molecular formula of C₂₇H₃₉N₆O₈P and a molecular weight of 602.6 g/mol. Remdesivir has the following structural formula:



11.1 Physical Appearance

Lyophilized Powder

Remdesivir for injection, 100 mg, is a sterile, preservative-free lyophilized powder that is to be reconstituted with 19 mL of Sterile Water for Injection and diluted into 0.9% saline prior to administration by intravenous infusion. Remdesivir for injection, 100 mg, is supplied in a single-dose clear glass vial.

The appearance of the lyophilized powder is white to off-white to yellow lyophilized powder or lumps or solid.

11.2 Inactive Ingredients

The inactive ingredients are sulfolbutylether-β-cyclodextrin sodium salt (SBECD), Water for Injection, USP, and may include hydrochloric acid and/or sodium hydroxide for pH adjustment. Remdesivir for injection, 100 mg, contains 3 g SBECD.

12. CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Remdesivir is an adenosine nucleotide prodrg that distributes into cells where it is metabolized to form the pharmacologically active nucleoside triphosphate metabolite. Metabolism of remdesivir to remdesivir triphosphate has been demonstrated in multiple cell types. Remdesivir triphosphate acts as an analog of adenosine triphosphate (ATP) and competes with the natural ATP substrate for incorporation into nascent RNA chains by the SARS-CoV-2 RNA-dependent RNA polymerase, which results in delayed chain termination during replication of the viral RNA. Remdesivir triphosphate is a weak inhibitor of mammalian DNA and RNA polymerases with low potential for mitochondrial toxicity.

12.2 Pharmacokinetics

The pharmacokinetics (PK) of remdesivir have been evaluated in adults in several Phase 1 trials.

- The pharmacokinetics of remdesivir and metabolites have not been in evaluated in patients with COVID-19
- Following single-dose, 2-hour IV administration of remdesivir solution formulation at doses ranging from 3 to 225 mg, remdesivir exhibited a linear PK profile.
- Following single-dose, 2-hour IV administration of remdesivir at doses of 75 and 150 mg, both the lyophilized and solution formulations provided comparable PK parameters (AUC₀₋₁₂, AUC₀₋₂₄, and C_{max}), indicating similar formulation performance.
- Remdesivir 75 mg lyophilized formulation administered IV over 30 minutes provided similar peripheral blood mononuclear cell (PBMC) exposure of the active triphosphate metabolite GS-443902 as remdesivir 150 mg lyophilized formulation administered IV over 2 hours.
- Following a single 150 mg intravenous dose of [¹⁴C]-remdesivir, mean total recovery of the dose was greater than 92%, consisting of approximately 74% and 18% recovered in urine and feces, respectively. The majority of remdesivir dose recovered in urine was metabolite GS-441524 (49%), while 10% was recovered as remdesivir.

Specific Populations

Sex, Race and Age

Pharmacokinetic differences based on sex, race, and age have not been evaluated.

Pediatric Patients

The pharmacokinetics of remdesivir in pediatric patients has not been evaluated.

Physiologically-based pharmacokinetic models were developed to estimate remdesivir and GS-441524 exposure and predict pediatric patient exposure based on age-dependent physiologic changes (e.g., organ volume/ function, blood flow). These simulations do not account for the impact of infection on the pharmacokinetics of remdesivir and GS-441524, which is currently unknown.

Renal Impairment

Because the excipient SBECD is renally cleared and accumulates in patients with decreased renal function, administration of drugs formulated with SBECD (such as remdesivir) is not recommended in adult and pediatric patients (>28 days old) with eGFR less than 30 mL per minute or in full-term neonates (≥7 days and ≤28 days old) with serum creatinine clearance ≥1 mg/dL unless the potential benefit outweighs the potential risk.

13. MICROBIOLOGY/RESISTANCE INFORMATION

Antiviral Activity

Remdesivir exhibited cell culture antiviral activity against a clinical isolate of SARS-CoV-2 in primary human airway epithelial (HAE) cells with a 50% effective concentration (EC₅₀) of 9.9 nM after 48 hours of treatment. The EC₅₀ values of remdesivir against SARS-CoV-2 in Vero cells was 137 nM at 24 hours and 750 nM at 48 hours post-treatment.

Resistance

No clinical data are available on the development of SARS-CoV-2 resistance to remdesivir. The cell culture development of SARS-CoV-2 resistance to remdesivir has not been assessed to date. Cell culture resistance profiling of remdesivir using the rodent CoV murine hepatitis virus identified 2 substitutions (F476L and V53L) in the viral RNAdependent RNA polymerase at residues conserved across CoVs that conferred a 5.6 fold reduced susceptibility to remdesivir. The mutant viruses showed reduced viral fitness in cell culture and introduction of the corresponding substitutions (F480L and V557L) into SARS-CoV resulted in 6-fold reduced susceptibility to remdesivir in cell culture and attenuated SARS-CoV pathogenesis in a mouse model.

14. NONCLINICAL TOXICOLOGY

The nonclinical toxicology profile of remdesivir has been characterized through the conduct of repeat-dose studies in rats and cynomolgus monkeys with once daily dosing up to 4 weeks in duration, studies to evaluate the genotoxic potential of the compound, a battery of reproduction and developmental studies (fertility in rats, embryofetal development in rats and rabbits, and a pre- and postdevelopmental study in rats), and a hemolysis/

blood compatibility study. Following repeated dosing in rats and monkeys, the kidney was identified as the target organ. In both species, clinical chemistry, urinalysis, and/or urinary biomarkers were early predictors of the observed kidney changes.

Carcinogenesis

Given the short-term administration of remdesivir for the treatment of COVID-19, long-term animal studies to evaluate the carcinogenic potential of remdesivir are not required.

Mutagenesis

Remdesivir was not genotoxic in a battery of assays, including bacterial mutagenicity, chromosome aberration using human peripheral blood lymphocytes, and *in vivo* rat micronucleus assays.

Impairment of Fertility

Nonclinical toxicity studies in rats demonstrated no adverse effect on male fertility at exposures of the predominant circulating metabolite (GS-441524) approximately 2 times the exposure in humans at the RHD. Reproductive toxicity, including decreases in corpora lutea, numbers of implantation sites, and viable embryos, was seen when remdesivir was administered intravenous daily at a systemically toxic dose (10 mg/kg) in female rats 14 days prior to mating and during conception; exposures of the predominant circulating metabolite (GS-441524) were 1.3 times the exposure in humans at the RHD.

Animal Toxicology and/or Pharmacology

Intravenous administration (slow bolus) of remdesivir to male rhesus monkeys at dosage levels of 5, 10, and 20 mg/kg/day for 7 days resulted, at all dose levels, in increased mean urea nitrogen and increased mean creatinine, renal tubular atrophy, and basophilia and casts.

Intravenous administration (slow bolus) of remdesivir to rats at dosage levels of ≥3 mg/kg/day for up to 4 weeks resulted in findings indicative of kidney injury and/or dysfunction.

15. ANIMAL PHARMACOLOGIC AND EFFICACY DATA

It is unknown, at present, how the observed antiviral activity of remdesivir in animal models of SARS-CoV-2 infection will translate into clinical efficacy in patients with symptomatic disease. Key attributes of the remdesivir nonclinical profile supporting its development for the treatment of COVID-19 are provided below:

- Remdesivir showed cell culture antiviral activity against a clinical isolate of SARS-CoV-2 in primary HAE cells (EC₅₀ value= 9.9 nM). The EC₅₀ values of remdesivir against SARS-CoV-2 in Vero cells has been reported to be 137 nM at 24 hours and 750 nM at 48 hours post-treatment.
- Remdesivir showed antiviral activity in SARS-CoV-2-infected rhesus monkeys. Administration of remdesivir at 10/5 mg/kg (10 mg/kg first dose, followed by 5 mg/kg once daily thereafter) using IV bolus injection initiated 12 hours post-inoculation with SARS-CoV-2 resulted in a reduction in clinical signs of respiratory disease, lung pathology and gross lung lesions, and lung viral RNA levels compared with vehicle-treated animals.

16. CLINICAL TRIAL RESULTS

Remdesivir is an unapproved antiviral drug with available data from two randomized clinical trials and a compassionate use program in patients with COVID-19, and from clinical trials in healthy volunteers and subjects with Ebola virus disease.

Clinical Trials in Subjects with COVID-19

NIAD ACTT-1 Study

A randomized, double-blind, placebo-control clinical trial evaluated remdesivir 200 mg once daily for 1 day followed by remdesivir 100 mg once daily for 9 days (for a total of up to 10 days of intravenously administered therapy) in hospitalized adult patients with COVID-19. The trial enrolled 1063 hospitalized patients in a 1:1 manner to receive remdesivir or placebo. The primary clinical endpoint was time to recovery within 28 days after randomization. In a preliminary analysis of the primary endpoint performed after 606 recoveries were attained, the median time to recovery was 11 days in the remdesivir group compared to 15 days in the placebo group (hazard ratio 1.31; 95% CI 1.12 to 1.54, p<0.001). Mortality was 8.0% for the remdesivir group versus 11.6% for the placebo group (p=0.059).

Study GS-US-540-5773

A randomized, open-label multi-center clinical trial (Study GS-US-540-5773) of patients with severe COVID-19 compared 197 adult patients who received remdesivir 200 mg once daily followed by remdesivir 100 mg once daily for 9 days (for a total of 10 days of intravenously administered therapy) with 200 adult patients who received remdesivir 200 mg once daily followed by remdesivir 100 mg for 4 days (for a total of 5 days of intravenously administered therapy), plus standard of care. The primary clinical endpoint was clinical status assessed by a 7-point ordinal scale at Day 14 after randomization. The study suggested that patients receiving a 10-day treatment course of remdesivir had similar improvement in clinical status compared with those receiving a 5-day treatment course (10-to-5 day odds ratio: 0.76; 95% confidence interval [CI] 0.51 to 1.13) on Day 14).

Clinical improvement was defined as an improvement of two or more points from baseline on a predefined 7-point scale, ranging from hospital discharge to increasing levels of oxygen support to death. Patients achieved clinical recovery if they no longer required oxygen support or were discharged from the hospital.

The time to clinical improvement for 50% of patients was 10 days in the 5-day treatment group and 11 days in the 10-day treatment group. At Day 14, observed rates between the 5- and 10-day treatment groups were 65% vs 54% for clinical improvement, 70% vs 59% for clinical recovery, and 8% vs 11% for mortality.

Compassionate Use Program in Patients with COVID-19

Remdesivir has been provided through a compassionate use multi-center, openlabel program to over 1,200 adult patients with confirmed SARS-CoV-2 infection by polymerase chain reaction (PCR) and manifestations of severe disease. In addition, remdesivir has been provided to 76 pediatric patients <18 years of age and 96 pregnant women through the compassionate use program.

Patients were treated with remdesivir 200 mg once daily followed by remdesivir 100 mg for 9 days intravenously, plus standard of care, for a total of up to 10 days of therapy.

Clinical Studies in Healthy Adults

Remdesivir was evaluated in four Phase 1 studies in 138 healthy adult volunteers (Studies GS-US-399-1812, GS-US-399-1954, GS-US-399-4231, and GS-US399-5505). In these studies, transient graded elevations in ALT and AST were observed at repeated once-daily doses of remdesivir.

Clinical Study in Subjects with Ebola Virus Disease

Supportive safety data are provided from the PALM study, a Phase 2/3, openlabel, randomized, parallel group study to assess the safety and efficacy of investigational treatments, including remdesivir, in patients with Ebola virus disease. 175 patients were randomized to receive remdesivir. A total of 9 SAEs judged by the site investigator as not related to underlying Ebola virus disease were reported for participants receiving remdesivir. Of these, an event of hypotension, which occurred during administration of the loading dose and led to fatal cardiac arrest, was considered related to remdesivir. The independent pharmacovigilance committee noted that the death could not be readily distinguished from underlying fulminant Ebola virus disease.

17. HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

Lyophilized Powder

Remdesivir for injection, 100 mg, is supplied as a single-dose vial containing a sterile, preservative-free White

to off-white to yellow lyophilized powder or lumps or solid that is to be reconstituted with 19 mL of Sterile Water for Injection and diluted into 0.9% saline prior to administration by intravenous infusion. Following reconstitution, each vial contains 5 mg/mL remdesivir reconcentrated solution with sufficient volume to allow withdrawal of 20 mL of 5 mg/mL solution containing 100 mg of remdesivir.

Discard unused portion.

The container closure is not made with natural rubber latex.

Storage and Handling

Do not reuse or save unused remdesivir lyophilized powder, for infusion for future use. This product contains no preservative.

Lyophilized Powder

Store remdesivir for injection, 100 mg, vials below 30°C (below 86°F) until required for use. Do not use after expiration date.

After reconstitution, vials can be stored up to 4 hours at room temperature (20°C to 25°C [68°F to 77°F]) prior to administration or 24 hours at refrigerated temperature (2°C to 8°C [36°F to 46°F]). Dilute within the same day as administration.

For further information write to: ProductSafety@mylan.com

POM | Schedule 2 | PP

To be distributed and used only in the licensed countries identified below:

Licensed Countries		
Afghanistan	Fiji	Nigeria
Algeria	Gabon	Pakistan
Angola	Gambia, The	Palau
Anguilla	Georgia	Panama
Antigua & Barbuda	Ghana	Papua New Guinea
Armenia	Grenada	Philippines
Aruba	Guatemala	Rwanda
Azerbaijan	Guinea	Samoa
Bahamas	Guinea Bis	Sao Tome and Principe
Bangladesh	Guyana	Senegal
Barbados	Haiti	Seychelles
Belarus	Honduras	Sierra Leone
Belize	India	Sint Maarten (Dutch part)
Benin	Indonesia	Solomon Islands
Bermuda	Jamaica	Somalia
Bhutan	Kazakhstan	South Africa
Botswana	Kenya	South Sudan
British Virgin Islands	Kiribati	Sri Lanka
Burkina Faso	Korea, Dem. People's Rep. (North Korea)	St. Kitts and Nevis
Burundi	Kyrgyzstan	St. Lucia
Cambodia	Lao DR (Laos)	St. Vincent and the Grenadines
Cameroon	Lesotho	Sudan
Cape Verde	Liberia	Suriname
Cayman Island	Libya	Tajikistan
Central Af R.	Madagascar	Tanzania
Chad	Malawi	Thailand
Comoros	Maldives	Timor-Leste
Congo DR	Mali	Togo
Congo Rep.	Marshall Islands	Tonga
Cook Islands	Mauritania	Trinidad & Tobago
Costa Rica	Mauritius	Tunisia
Cote d'Ivoire	Micronesia, Fed. Sts.	Turkmenistan
Cuba	Moldova	Turks & Caicos
Curacao	Mongolia	Tuvalu
Djibouti	Montserrat	Uganda
Dominica	Morocco	Ukraine
Dominican Republic	Mozambique	Uzbekistan
Egypt	Myanmar	Vanuatu
El Salvador	Namibia	Vietnam
Equatorial Guinea	Nauru	Zambia
Eritrea	Nepal	Zimbabwe
Eswatini (Swaziland)	Nicaragua	
Ethiopia	Niger	



Manufactured by:
Mylan Laboratories Limited
At: Plot No. 42 to 52, Survey No. 166, 171, 172 & 177,
TSIIC, Phase -II, IDA, Pashamylaram (V), Patancheru (M),
Sangareddy District, Telangana State – 502307.

Code No.: TS/DRUGS/SGY/2020-65910
TM - Trade Mark under registration

DESREM is manufactured under a license from Gilead Sciences, Inc.



0108

ARTWORK DETAIL LABEL

Product	DESREM - Remdesivir for Injection 100 mg/vial		
Buyer/Country	Mylan / ROW (General Export) - (GLAND)	Component	Pack Insert
Dimension	420 x 290 mm	Pack	--
New Item Code	1032158	Old Item Code	NA
Colour Shades	Black	No. of Colours	1

Change Control No.	NA
Design/Style	Front & Back Printing. To be supplied in folded size of 55 x 35 mm. Brand name facing front after final folding.
Substrate	40/45 GSM Paper.
Special Instructions	Printing clarity to be clear & sharp.
Autocartanator Requirements	NA
Caution to the printer: Before processing, please ensure that the ARTWORK received for printing is exactly in line with APPROVED ARTWORK provided to you. In case of any FONTS/DESIGN are Mis-matching with the APPROVED ARTWORK, please inform PDC for further action. DO NOT MAKE ANY CHANGE TO THE ARTWORK WITHOUT WRITTEN INSTRUCTIONS FROM PDC.	

SOP - 000484829

FORM - 000589186 V 1.0

Rev. 0; Ver. 2; Date: 02-10-2020

BACK SIDE