Georgia HCV Elimination Spring Workshop

7 & 9 March 2018

Venue: National Center for Disease Control and Public Health (NCDC), Tbilisi, Georgia

Participants: Approx. 100

SUMMARY

- Organized by: The Georgia Ministry of Labour, Health, and Social Affairs (MoLHSA) and the Georgia NCDC.
- Objective: To bring local stakeholders and program implementers together to discuss program progress, successes, and challenges and to review TAG recommendations and how best to implement the recommendations.
- Challenges remain to reach the 2020 HCV elimination goals. Decentralization of screening, care, and treatment services will play a key role in achieving the goals and Georgia is investing greatly in this strategy.
- Diversity of services is critical: Although decentralization will play a key role in reaching the goals, other service models, i.e. the existing treatment centers will continue to play an important role in guiding care and treatment services nationally and in providing care to more severely ill patients.
- Decentralization workgroups: Overall, Primary Care/Hospital, and Harm Reduction will play a critical role.
 Key tasks include:
 - Workgroups meet monthly and urgently develop implementation plans
 - o Ensure quality monitoring and evaluation/data collection
 - Ensure lessons learned from the model/pilot programs are considered when developing National implementation plans
 - Simplified care and treatment regimens are critical. Pangenotypic regimens will play a major role in simplification
 - o Partnerships will be critical to successful National implementation
- Infection prevention and control in healthcare settings is making progress and there may be opportunities to collaborate with other efforts.
- Research efforts will continue to inform the National Program, both in primary care and in harm reduction
- A need to develop policies for testing for re-infection in harm reduction settings. This requires viral core
 antigen or PCR testing and is not currently covered by the program. It was recommended that a free of
 charge annual test could be included for beneficiaries.
- Decreasing time from screening to confirmation to returning results could improve linkage to care. Efforts should be made to decrease this timeframe. Making services available to all who are confirmed with infection without committee review was recommended as a solution.
- It was recognized that drug security can be deemphasized, as drug prices have dropped greatly throughout the world.
- Information technology needs to be responsive to the program and incorporate decentralized services.
- Laboratory services need to be expanded to accommodate decentralized services.
- Decentralization of services needs to be sensitive to potential unanticipated costs such as the need for doctors in harm reduction sites.
- Decentralization to harm reduction sites could lead to more accurate data regarding numbers of PWID receiving services; however, ensuring confidentiality remains a significant barrier.
- To the degree possible, co-locating or coordinating care and treatment services between harm reduction and primary healthcare could alleviate costs and other barriers.

The 5th National Hepatitis C Elimination Workshop Day 1 – March 7

Time	Topic	Speaker	
09:30-10:00	Welcome/Introduction	David Sergeenko, Minister of	
09.50-10.00	welcome/introduction	Labour, Health and Social affairs	
		Elisabeth Rood, Deputy Chief of	
		Mission, Embassy of the US in	
		Georgia	
		Dr. Beth Skaggs, CDC	
10:00-10:15	Media Coverage	DI. Detil Skaggs, CDC	
10:15-10:40	Review Progress of Elimination Program	Amiran Gamkrelidze, NCDC	
10:40-11:00	Brief overview of agenda and purpose of the meeting	Francisco Averhoff, CDC	
11:00-11:30	Coffee Break	Trancisco Avernori, ese	
11.00 11.50	Decentralization HCV treatment access through		
	decentralization of screening, care and treatment services		
11:30-11:50	Protocol of implementation	Eka Adamia, MoLHSA	
11:50-12:00	Lessons learned from HIV scale up	Philippa Easterbrook, WHO	
	'		
12:00-13:00	Guided Discussion to develop plan for way forward	MOLHSA	
12.22.11.22	timeline for activities	Francisco Averhoff, CDC	
13:00-14:00	Lunch		
	Introduction and progress update from Primary Care and		
	Hospitals Decentralization Workgroup		
14:00-14:20	a lateralization and progress undetections Drives	Tengiz Tsertsvadze, Infectious	
14.00-14.20	 Introduction and progress update from Primary care and Hospitals W/G 	Diseases, AIDS and Clinical	
	care and nospitals w/O	Immunology Research Center	
	Updates on projects	(IDACIRC)	
14:20-14:35	Integrating HCV screening and simplified treatment	Tengiz Tsertsvadze, IDACIRC	
	services in primary healthcare		
14:35-14:50	Simplification of pretreatment diagnostic	Jaba Zarqua, Mrcheveli	
	evaluation and on-treatment monitoring	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	
	procedures within HCV Elimination Project		
	(Mrcheveli)		
14:50-15:05	HIV, TB, and HCV at primary healthcare settings—A	Irma Khonelidze, NCDC	
	pilot project in Samegrelo-Zemo Stvaneti		
15:05-16:05	Guided Discussion to develop plan for way forward timeline	Tengiz Tsertsvadze, IDACIRC	
	for activities ,	-	
16:05-16:20	Coffee break		
	Infection Control		
	Improving IPC in Georgia, Future activities, gaps		
16:20-16:30	MOLHSA Activities	Marina Darakhvelidze, MoLHSA	
16:30-16:40	NCDC Activities	Giorgi Chakhunashvili, NCDC	
16:40-16:50	CDC Activities	Meghan Lyman, CDC	
16:50-17:30	Guided Discussion to develop plan for way forward timeline	Marina Darakhvelidze, MoLHSA	
	for activities	Paata Imnadze, NCDC	
17:30-18:00	WHO guidance on viral hepatitis surveillance	Antons Mozalevskis, WHO	
18:00-18:15	Closing Remarks	Francisco Averhoff, CDC	

Day 2 – March 9

Time	Topic	Speaker
09:30-09:45	Opening/Welcome remarks	Francisco Averhoff, CDC
09:45-10:05	Integration of HCV treatment Services in Harm reduction Centers • Introduction and progress update from Harm Reduction Decentralization Workgroup Updates on projects	Maia Butsashvili, HRU
10:05-10:20	 Implementing HCV treatment in harm reduction centers in Georgia (HRU Project) 	Maia Butsashvili HRU
10:20-10:35	FIND "HEAD-Start project in Georgia"	Francois Lamoury, FIND
10:35-10:50	Barrier Study in HR Centers	Maka Gogia, Harm Reduction
10:50-11:10	Coffee Break	
11:10-12:20	Guided Discussion to develop plan for way forward timeline for activities	Francisco Averhoff, CDC Irma Khonelidze, NCDC
12:20-13:20	Lunch	
13:20-13:40	 HCV Laboratory Diagnostics Quality of HCV testing in Georgia through National External Quality Assurance Program 	Dr. Beth Skaggs, CDC
13:40-14:00	Confirmation of HCV infection by Core Ag using a centralized laboratory approach	Dr. Nazi Chitadze, NCDC
14:00-14:25	 Simplified diagnostics and treatment monitoring of HCV infected patients 	Karla Thornton, Project ECHO
14:25-15:25	Guided Discussion to develop plan for way	Beth Skaggs, CDC
	forward timeline for activities	Maia Alkhazashvili, NCDC
15:25-15:40	Coffee Break	
15:40-16:00	Scientific Committee Progress	Tatia Kuchuloria, CDC
16:00-16:15	 Introduction of Cured Hepatitis C Patients' Association 	Temur Radiani, Cured Hepatitis C Patients' Association
16:15-16:40	Blood Safety Status of blood safety program in blood banks, successes, challenges, and plans to address challenges	Eter Kipiani, NCDC
16:40-17:30	 Guided Discussion to develop plan for way forward timeline for activities 	Marina Darakhvelidze, MoLHSA Khvicha Getia, NCDC
17:30-18:00	Closing Remarks	David Sergeenko Francisco Averhoff



Decentralisation of HCV testing, care and treatment: Lessons learned from HIV

Dr. Philippa Easterbrook

Global Hepatitis Programme

and HIV Department



5th National Hepatitis C
Elimination Workshop
Tbilisi, Georgia
7-9 March 2018



Outline of presentation

- HIV and HCV: Similarities and differences
 - Common barriers to increasing treatment access
 - Common features and differences epidemiology and treatment approach
- Progress in global hepatitis C and HIV response
- Key lessons learnt from HIV/ART scale-up
 - Testing
 - Linkage to Care
 - Treatment and Service Delivery
- What is next?

Common initial barriers to increasing testing and treatment access


Barrier	HIV	HCV	HBV
Limited data on epidemiological situation	✓	✓	✓
High stigma and discrimination	✓	✓	✓
Lack of political and financial commitment	✓	✓	✓
Complexity of drug regimens, high pill burden Side effects that complicate adherence	✓	✓	
Complexity of screening, treatment and follow-up monitoring	✓	√	✓
High cost of drugs and tests	✓	✓	✓
Patenting of drugs approved and in pipeline	√	✓	✓
Highly specialized vertical services	✓	✓	✓
Limited patient and community engagement	√	√	√

Similar, yet different.... Epidemiology and Treatment

	2016/2015	HIV	HCV
	Main transmission routes	Sexual IDU	Unsafe healthcare injections IDU
	Burden of infection	36.7 million (HIV Ab)	71 m (HCV RNA)
	Deaths per year	1 million	399,000
	New infections per year	1.8 million	1.75 million
	% infected requiring treatment	Treat All All HIV Ab positive	Treat All All HCV RNA positive
	Treatment duration	Lifelong	12 weeks
	Treatment response	70-80% HIV RNA undetectable (1yr)	>95% Cure HCV RNA undetectable (SVR12)
	Numbers treated	19.5 million	3 million
	% diagnosed/%treated	70%/53%	20%/7%
4	Funding	Global Fund/PEPFAR \$19 billion per year	Domestic

Progress in ART and DAA scale-up

ART scale up over the last decade



Persons starting curative HCV treatment with DAAs

2014: < 200 000


2015: 1.1 million

2016: 1.7 million


TOTAL: ~ 3 million

Source: WHO, work conducted by CDA/Polaris

Eliminate viral hepatitis as a major public health threat by 2030, as defined by:



Global elimination 2030 targets and Where are we at now?



Coverage (%)

2015 baseline

2030 targets

(A)

Key lessons learnt from HIV/ART scale-up

- I. The "Public health approach" to HIV care and treatment
- II. Strategies to reduce drug costs and increase access
- III. Success is possible: increasing HIV testing access/uptake
- IV. Where to test? Learning from evolution in HIV testing approaches
- V. Simplified diagnostic and management algorithms
- VI. Linkage to Care is key
- VII. Simplified drug regimens and WHO treatment guidelines
- VIII. Evolution in "when to start" ART to "treat all" in HIV care
- IX. Service delivery: Decentralization of testing + treatment and Task shifting to support decentralization
- X. Differentiated care
- XI. Integration
- XII. Tracking progress: the cascade of care

Lessons on Expanded Access to Testing

Viral hepatitis testing is at core of entry to treatment and prevention cascade

LESSON 4

Where to test? Learning from evolution in HIV testing approaches

	TESTING APPROACHES	
TESTING SITES	Routinely	Focused (Risk-base
HEALTHCARE FACILITY		1 2 2
TESTING		
Primary care settings	X	
Antenatal clinics	×	
HIV clinics	×	
TB clinics	×	
STi clinics	×	1
Drug treatment and harm reduction services	×	
Inpatient and outpatient hospital settings	×	
Paediatric and adolescent clinics	X	
COMMUNITY-BASED		1
Mobile/outreach testing for priority populations		×
Mobile/outreach for the general population (for example young people)		×
National testing campaigns/camps	х	×
Testing of family members		X
Partner testing (for all partners of people with viral hepatitis)		×
Mass media and social media	×	×
Home-based/ door-to-door testing	х	×
Workplace testing		×
School/educational institution testing		×




Health provider Focussed testing in health facilities/outreach

- High acceptance (HIV, antenatal & TB)
- But still many missed opportunities for testing
- Partner testing


Moving Testing Out of Health Center into community

- Home-based (house to house), Campaigns, Outreach (mobile), partner, workplace
- Lay provider testing (already policy in 64% of SSA countries)
 - Self-testing



General pop testing: Egypt (village based programme), Mongolia and Georgia (PHC)

LESSON 5 Simplified testing and management algorithm



Lessons on promoting Linkage to Care

High rate of loss to follow up after positive serology, and low uptake of HCV RNA and access to treatment


LESSON 6

Linkage to care is key (HIV and HCV)

Strategies to consider for improving linkage

Evidence is limited, but promising

- Integration of services (testing/treatment) at a single facility, especially within mental health/drug treatment services
- On-site or immediate RDT testing with same day results
- Trained Peer and lay health worker support to identify and reach people lost to follow-up
- Clinician reminders to prompt provider initiated, facility-based testing plus reflex NAT
- 13 DBS specimens for NAT ±serology in some settings



For the future...

Decentralised viraemic testing and treatment

Lessons on Expanded Access to Treatment

LESSON 8a: - Evidence for clinical impact Starting earlier reduces mortality and morbidity

The NEW ENGLAND JOURNAL of MEDICINE

20104112020 CH 1011

AUGUST 27, 2015

ma. 272 340. 6

Initiation of Antiretroviral Therapy in Early Asymptomatic HIV Infection

The DUSIGHT START Study Groups

ARRESTS ACT

商金 に見込まの日本日

Data from randomized stude are backing in the bunelin and rios of including authorities of therapy in passens with apstigue-stude human intrinsusded intersion (LHV) infection when have a CD4+ cours of more class 250 cells per cubic millianum.

efficant

We transforth assigned HPS-positive adults who had a CD4+, count of more than 500 cells per cable millionner to sum adirectorical sheapy homeditanly disministration appears to the substance of the count decreased to 550 cells per cable millioners of until the development of the sugained intimanodericonses acidemic AULS; or another conditions, that distant the use of automorphism develop inference-instants group. The primary composite and point with any serious AULS-randed overs, artiment man-AULS-related overs, or death from any case.

Below of B

A moral of 4465 panisms were liferweed for a means of 3.0 years. As souly story, the median HIV visit load was \$12,799 colored per infelligence, and the triedlan CO4-count wiso 651 calls per cubic millimeter. On May 15, 2015, on the basis of an insertion analysis, the disas and safety mosacieties, possible, con the basis of an insertion analysis, the disas and safety mosacieties produce the extraordinate special per strong per strong analysis, and the production of the parisms in the desirent-printing action property and the strong per strong per strong analysis of the period period

COMPLETEDIORS

The Instantin of antirenrosins' therage in 18T9-possible adults with a GC-4- count of more than 500 cells per cube millimener photoded into benefits over starting such through in patients after the CG-4- count had declined in 250 cells per cubic millitation. (Studed by the National Institute of Allengs and Institutes Diseases and orbane STARK ChicaFiblia.gov sumber. 18CC00867-044;


the newsless of the newsless group lives to Lordgens, M.D., Questiany Scholt 14, Stables, M.D., Stables, Scholt Stables, M.D., Stables, S. Seel Stables, M.D., Stables, S. Seel Stables, M.D., Garell, M.D., Stables, M.D., Garell, M.D., Stables, M.D

No complete list of secretary in the Disategic Taxing of districtionical Transforms (2 NAT) Study Umage is provided in the longitumismum Approach, as a fable of NAM copy.

This as to be very published to the july 26, 2007.

This article was published on judy on 2005 at 1975 at 1975.

IN COURT OF THE MANAGEMENT OF



LESSON 8b: - Evidence for public health impact HIV viral suppression reduces incidence

Community viral load, antiretroviral therapy coverage, and HIV incidence in India: a cross-sectional, comparative study



Send Solver Science, Shoot HW etc., Allow M McFall, Aglar K Selt inhous, Shorm your Streaming, Olive Lansended et. Psychorouthy Buldrighner, David D Gillertung Scriet Schoron Green Williams

Exchanged HIV Incidence is the best measure of grammen-programme effects once; but his measurement is transfered and a subdifficult and exponence. The concept of community vital load as a modifiable driver of new HIV intections has assumed because of the subscantial amendion. We see our to compare several measures of community viral load and antinensyiral therapy. Markot make (ART) coverage as correlates of HTV incidence in high-risk populations.

Mediods We analysed data from a sample of people who inject drugs and men who have sex with men, who were participants of the baseline assessment of a cluster-randomised trial in progress across 22 cities in India (Clinical trials gov number NCT05686750). We recruised the study population by use of respondent-driven sampling modern theorem at the and did the baseline assessment at 27 community-based sites (12 for men who have see with men and 15 for becole who inters drags). We estimated HIV incidence with a multisony algorithm and calculated five community-based measures of HIV control mean log., HIV RNA in participants with HIV in a community either engaged in care (In-care viral lead), aware of their status but not necessarily in care (aware viral lead), or all HIV positive individuals must be presented whother they were aware, in care, or not (notulation viral load); participants with HIV in a community with HIV ENA Spokening, heaven, at more than 150 copies per ml. (prevalence of viraemia); and the proportion of panicipants with HIV who self-reported ART use in the previous 30 days (population ART coverage). All participants were usted for HIV, with additional using in HIV-positive individuals. We assessed correlations between the measures and HIV incidence with Spearman correlation coefficients and linear regression analysis.

Findings Between Oct 1, 2012, and Dec 19, 2013, we recruised 26 503 participants, 12022 men who have six with men and A Colombia DA 14.451 people who inject drugs. Median incidence of HIV was 0.57% (IQR 0.40-1-17) in men who have see with men. Harmon Po. and 1.43% (0.60-4.00) in people who inject drugs. Prevalence of virannia was more strongly correlated with HIV Incidence (correlation 0-81, 95% CI 0-62-0-91; p-0-0001) than all other measures, although correlation was significant. with aware viral load (0.59, 0.27-0.79; p=0.001), population viral load (0.51, 0.1s-0.74; p=0.007), and population ART coverage (-0.54, -0.76 to -0.20; p=0.000). In carry-frail load was not correlated with HIV incidence (0.29, -0.10 to 0.00. p=0.14). With regression analysis, we estimated that to reduce HIV incidence by 1 percentage point in a community. Section 80,150 prevalence of virarmia would need to be reduced by 4-34%, and ART use in HIV-positive individuals would need to

Interpretation Providence of viraentia had the serongest correlation with HIV incidence in this sample and might be a useful measure of the effectiveness of a measurem programme.


Funding US National Institutes of Health, Elson John AIDS Foundation

Compelling evidence of HIV examine as presented from clinical southes' and expending but highly variable giobal antimerowiral thorapy (ART) coverage²³ has incured data derived from administrative databases do not assention on the need so monitor community-lovel effectiveness of ART. HTV incidence is the best possible mogram of the officebours of treatment as provention account for a large proportion of HIV transmission. programmes. However, measurement of HIV incidence is Community viral load measured in the consest of a difficult and exponents, and rarely done outside a research cohort study might represent individuals out of care, but contact." Researchers have proposed the measures of it is unlikely to adequately account for individuals who community viral loads and population ART coverage as are unaware of their status, because seronisgative cohercapturing a causal association between community level participants are usually routinely ested for HTV at study officety-chass of HIV grounding and HIV gransmission.

individuals with HIV, has been temporally correlated which can vary widely by community

with HTV incidence or new HTV diagnoses (from adminiscrative databases) in emingical sending.18 However, these measures have limitations. Community viral lead represent individuals who are out of care or unaware of their status a population that has been proposed to visies. Some researchers have argued that omeral Community viral load, a family of measures derived sendency measures of community viral load among from HIV RNA measurements in subgroups of poorle with HIV do not account for HIV prevalence.


linkers largest ten School of Medicine, Department of CW Lucus MCh. Johnn Hopki Short D.D.Dalamograph Scill Secretary and Relatedors Plaintrolman Phil National Institute of Allerton Or Suppl Subject Solomon intern Hookins School o



LESSON 8c: - Evidence for programmatic impact High rate of loss to follow up if not treated



New studies show that ART can be started on same day as HIV diagnosis



WHO recommendation (July 2017)

Start within 7 days of an HIV diagnosis

Consider same day start

•

PLOS Medicine I DOI:10.1371/journal.pmed.1002015 May 10, 2016

Lessons on Service Delivery approaches to expand access to testing and treatment

Simplified HCV Service Delivery for a Public Health Approach

Good service delivery health practice principles

- 1. Simplified and standardized algorithms
- 2. Strengthen Linkage from testing to care, treatment and adherence
- **3. Integrated** testing, care and treatment
- **4. Decentralisation of care** to promote access
- Differentiated care to address diverse health care needs of those with HCV
- 6. Engagement with Community and peer support

Service delivery in specific populations

- Persons who inject drugs
- People in prisons and other closed settings
- MSM and sex workers
- Adolescents and Children
- Migrant/indigenous populations
- Pregnant women






7. More efficient procurement + supply management

LESSON 9

Decentralization of testing and treatment supported by Task shifting

ART initiation and prescribing by nurses can be done safely, and improve health outcomes and quality of care



Time to ART initiation decreased from nearly 100 days in 2003 to less than 3 weeks in 2009





Figure 2 Access to antiretroviral therapy in Thyolo district; MA, medical assistant.


Bemelmans et al, Trop Med Int Health 2010

LESSON 10: - Differentiated care (based on evolving HIV care and treatment principles)




Who? HCV infected persons category	What? Care needs	Where? Site	By Whom? Care-giver
Persons clinically well and stable	Standard care package: Counselling, adherence support, treatment initiation and monitoring	Facility-based including primary care or community-based settings, including mobile/outreach	Physician or ?nurse
Advanced liver disease or serious co- morbidities, HCC, previous treatment failure	Requiring more intensive clinical support and follow-up: Management of liver related complications (eg. variceal bleed, ascites, encephalopathy, HCC treatment, genotyping)	Facility-based - hospital	Physician
Mental health issues, active injecting drug users, alcohol misuse, adolescents	Requiring more intensive psychosocial/mental health support	Can be Facility- based or Community- based, Harm reduction site	Physician and counsellor/peer support

LESSON 11 Integration




LESSON 12

Tracking progress: the cascade of care



Cascade of care for HCV infection by WHO region, 2016





Next steps

- What more can we learn from HIV?
 - Differentiated care, decentralization and task-shifting approaches
 - Self-testing
- What can HIV learn from hepatitis?
 - Demonstration projects and operational research on different models of care
- How can we integrate better?
 - When and how to share technologies

Thank you + Acknowledgements

Global Hepatitis Programme

Nathan Ford

Marco Vitoria

Meg Doherty

Michel Beusenberg

Rachel Baggaley

The 5th National Hepatitis C Elimination Workshop Tbilisi, Georgia

Introduction and Progress Update from Primary Care and Hospitals Working Group


Tengiz Tsertsvadze MD, PhD

Director General, Infectious Diseases, AIDS and Clinical Immunology Research Center

Professor, Tbilisi State University

Georgia Hepatitis C Elimination Program Care Cascade

April 28, 2015 - October 31, 2017



Source: Georgia's HCV Elimination Program Treatment Database

Rationale

- Increasing HCV treatment uptake is needed to achieve ambitious elimination goal
- Integrating screening and simplified treatment services in primary healthcare offers opportunity to eliminate barriers and to improve engagement in the entire continuum of HCV care
- Integrated and simplified models of care are effective in ensuring high diagnosis, treatment and large number of persons achieving SVR

First Georgia Meeting on Decentralization held at MoLHSA on December 8, 2017

Primary care /hospitals working group established


Objective

Develop a practical model of care to be implemented for HCV screening and treatment in primary health care centers and district hospitals that maximizes HCV screening, enrollment, and retention in the HCV program of the general population.

Roles and Responsibilities of Primary healthcare /hospitals working group

- Meet as a group as recommended by the oversight committee
- Adhere to timeline of task completion
- Selection of primary healthcare facilities across the country
- Develop an action plan for decentralization at PHCs and district hospitals that addresses the challenges
- Adhere to expectations of meetings, communication, documents/products set by the oversight workgroup

Decentralization of HCV treatment and care at PHCs



Major challenges

- Simplified Treatment and Diagnostics guidelines/algorithms
 - Necessary tests for screening; confirmatory testing; simplified pre-treatment; monitoring and follow-up diagnostics; location of testing; consolidation of testing
 - National standardized treatment guidelines; standardized medication supply chain
- Ensure laboratory and diagnostics integration
 - screening; confirmation; pre-treatment; monitoring and follow-up diagnostics; standardized interpretation of results
- Develop and disseminate SOPs
 - using simplified diagnostics plans and simplified treatment regimens

Major challenges (cont.)

- Develop Curriculum for PHC/hospitals
 - curriculums/training in screening, confirmation, pre-treatment, treatment and monitoring diagnostics
- Initial and Continued training (CME) for PHC and Hospital staff
 - areas of potential initial training; continued education/support/mentorship; in-person and teleconference modules
- Use of ECHO platform to facilitate training
 - curriculum development; initial training; continued education

Major challenges (cont.)

- Develop necessary information systems/modules
 - method of data collection and storage; integration into national Elim-C database; standardized data collection
- Elaborate the concept of treatment in rural areas (i.e. 69 districts)
- Incorporate lessons, findings from PHC/hospital related pilot projects into practical model/action plan





Pilot project

Integrating HCV screening and simplified treatment services in primary healthcare

This model will provide the basis for the decentralization of treatment and care in PHCs and hospitals nationwide

Completed activities

- A special accredited CME training program and curriculum for PHCs created initially for the pilot project to support training of primary care/hospitals providers countrywide
- Simplified HCV diagnostics and treatment protocols developed for the pilot project to be applied for primary care on national level
- Established ECHO telemedicine model to train and provide primary care providers and hospital staff with advice and clinical mentoring

Timeline of activities

- Simplified HCV diagnostics and treatment guideline/algorithms developed by April, 2018
- Primary care providers and hospital specialists under accredited CME training program are trained countrywide by June 2018
- ECHO telemedicine platform is expanded as a national model for using telemedicine for HCV treatment and care in rural areas by June 2018
- Laboratory diagnostics SOPs are develop and disseminated by June 2018

The 5th National Hepatitis C Elimination Workshop Tbilisi, Georgia

Integrating HCV screening and simplified treatment services in primary healthcare

Tengiz Tsertsvadze MD, PhD

Director General, Infectious Diseases, AIDS and Clinical
Immunology Research Center

Professor, Tbilisi State University





Pilot project

Integrating HCV screening and simplified treatment services in primary healthcare

Objectives/specific aims

To assess effectiveness of integrated HCV screening and treatment model in primary healthcare settings

Specific aims:


- Quantify engagement in the HCV care continuum (cascade) from screening through cure
- Assess efficacy and safety of simplified diagnostics and treatment monitoring procedures in primary healthcare settings
- Conduct cost-analysis of achieving sustained virologic response (SVR) with simplified approach and current standard of care
- Compare key project monitoring and evaluation indicators with national data

Methods

- Prospective evaluation of integrated screening and treatment program in PHCs
- Total of 4 PHCs will be selected, including one in Tbilisi and three in nonurban regional locations
- HCV screening and care services using "One stop shop" approach
- Patients will be offered anti-HCV test using an opt-out model. Same specimen will be used for confirmatory analysis
- Paperwork and registration with the MoLHSA will be completed by case managers at PHCs.
- Registered patients will be treated by primary care providers under the supervision of IDACIRC and HEPA clinic.

Pre-treatment evaluation

Algorithm for pre-treatment evaluation



Monitoring during antiviral therapy

Treatment monitoring approach will be simplified for PHCs

Simplified treatment monitoring procedures												
Measurements		ent Durati veeks)	on	After treatment completion (weeks)								
	4	8	12	12 or 24								
Clinical assessment	X	Х	X	X								
HCV RNA quantitative				X								
Complete blood count	X*	X*	X*									
ALT	X	X	Х									

^{*} only for patients receiving Ribavirin containing regimens

Guidance/supervision on integrated model implementation

IDACIRC and HEPA clinic together with other partners will provide guidance and supervision on integrated model implementation, including:

- Training on PHC personnel on hepatitis C related issues using
- Training on HCV rapid antibody testing
- Training on information system
- Utilizing the ECHO telemedicine model to train and support primary healthcare providers.
- Establishing M&E framework for assessing program effectiveness.

Completed activities



Completed activities (cont.)







- A special CME training program for primary care physicians developed and already approved by MoLHSA
- First training was conducted at the Family Medicine Training center in Tbilisi on March 5th.


Completed activities (cont.)

- HCV diagnostics, treatment and care protocol specially prepared for this project.
- Integration of the project activities into national HCV elimination program will be approved by a Government decree in nearest future.
- ➤ IT unit at MoLHSA has started to create a separate module in national HCV elimination program database to collect data on patients receiving diagnostics, treatment and care services at PHCs within this pilot project.

Future plans

- > Training of PHC personnel will continue
- Implementation of integrated model of HCV diagnostics and simplified treatment in PHC
- Implementation of ECHO telemedicine

Decentralization of HCV treatment and care at PHCs



Acknowledgements





John Ward, Francisco Averhoff, Muazzam Nasrullah Juliette Morgan, Beth Skaggs, Lia Gvinjilia, Shaun Shadaker, Tatia Kuchuloria, Archil Kublashvili



John Martin, Greg Alton, Graeme Robertson



Nezam Afdhal, Stefan Zeuzem, Geoffrey Dusheiko



Sanjeev Arora, Karla Thornton



David Sergeenko, Valeri Kvaratskhelia, Multisectoral Commission on Hepatitis C



Amiran Gamkrelidze

All HCV Clinical Care Provider Clinics

Simplification of pretreatment diagnostic and on-treatment monitoring procedures within HCV Elimination Project

The 5th National Hepatitis C Elimination Workshop
Tbilisi, Georgia, 2018

Dr. Jaba Zarkua, Medical Center Mrcheveli





Background

- From June 2016 Treatment for all HCV infected patients!
 - Main type of patients w/o advance liver disease.
 - Drugs available Sofosbuvir/Ledipasvir (Harvoni) +/- Ribavirin.
 - Treatment duration 12 or 24 weeks.
- Main problem decreasing number of patients.
 - Screening activities (challenge for 2017, activities are on-going)
 - Linkage to care (challenge for 2018)
 - Costs
 - Low motivation of patients
 - Low level of awareness
 - Disinformation about new drugs
 - Many mandatory procedures within project, discomfort

Standard of care procedures

- 1 Step: anti-HCV(+) patient's visit to Hub, perform confirmatory test (HCV RNA PCR/HCV core AG).
- 2 Step: Patient's first appointment to physician for diagnostics. Fasting state mandatory for lab. Investigations (glucose, ALP).
- 3 Step: Second visit to physician with lab. and imaging results. Final consultation before treatment. Complete clinical records and referral to service center.
- 4 Step: Start of treatment. Visit in every 2 weeks for drug control.
 Monthly for lab. tests.
- 5 Step: Follow up visit after 12 weeks from completing the treatment.

12 week regimens - schedule

Procedures	Pretreatment	2	4	6	8	10	12	Follow Up
Drug Control		Χ	Χ	Χ	Χ	Χ	Χ	
Physicians	XX		Χ		Χ		Χ	X
Consultation								
HCV RNA PCR	X		Χ					Χ
HCV Genotype	X							
CBC	X		Χ		X*		Χ	
ALT	Χ		Χ		Χ		Χ	
AST	Χ							
Creatinine	Χ						Χ	
Bil (tot., dir.)	X						Χ	
Albumin	X							
PT, INR	X							
ALP	Χ							
GGT	X							
Glucose (f.)	Χ							
HBsAg, anti-HBc	X							
(tot)								
Ultrasound	Χ							
Fibroscan	If needed							

^{*} RBV containing regimens only


24 week regimens - schedule

Procedures	Pretreatment	2	4	6	8	10	12	14	16	18	20	22	24	Follow Up
Drug Control		X	X	Χ	X	X	Χ	X	Χ	Χ	X	X	X	
Dharisiana	W		V		V		V						V	V
Physicians Consultation	XX		Χ		Χ		Χ						X	X
HCV RNA PCR	Х		X											X
HCV Genotype	X													
CBC	X		Χ		X*		Χ						Χ	
ALT	Х		Χ		Χ		Χ		X**		X**		Χ	
AST	X													
Creatinine	Х						Χ		X*		X*		Χ	
Bil (tot., dir.)	X						Χ						Χ	
Albumin	Х													
PT, INR	Х													
ALP	Х													
GGT	Х													
Glucose (f.)	Х													
HBsAg, anti-HBc (tot)	Х													
Ultrasound	Х													
Fibroscan	If needed													

^{*} RBV containing regimens only

^{**} RBV free regiments only

Pilot study by Mrcheveli/Neolab



Aims of the project

Primary aims:

- Reduce the patient's unnecessary clinical visits.
- Reduce the patient's unnecessary visits to HCV Elimination Service Center.
- Remove the unnecessary laboratory investigations.
- Reduce the financial expanses per patient.
- Reduce the budget of the Project.

Secondary aims:

- Simplified algorithm should not influence on the treatment outcome.
- Decrease the lost to follow up rates.
- Make participation in HCV Elimination Project more comfortable.

12 week regimens - Simplified

Procedures	Pretreatment	2	4	6	8	10	12	Follow Up
Drug Control		×	X	*	Χ	×	X	
Physicians Consultation	XX		X		X		X	X
HCV RNA PCR/HCVcAg	Χ		×					X
HCV Genotype	Χ							
СВС	Χ		×		×		×	
ALT	Χ		Χ		X		Χ	
AST	Χ							
Creatinine	Χ						×	
Bil (tot.)	Χ						×	
Dil (din)	X							
Albumin	Χ							
PT, INR	Χ							
ALD	X							
GGT	X							
Clusess (f.)	X							
HBsAg, anti-HBc (tot)	Χ							
Ultrasound	Χ							
Fibroscan	If needed							

24 week regimens - Simplified

Procedures	Pretreatment	2	4	6	8	10	12	14	16	18	20	22	24	Follow Up
Drug Control		×	Х	×	X	×	X	×	X	×	X	×	X	
Physicians Consultation	XX		X		X		X						X	X
HCV RNA PCR/HCVcAg	Х		×											X
HCV Genotype	Х													
СВС	Χ		X		×		X						×	
ALT	Χ		Χ		×		Χ		*		X		X	
AST	Χ													
Creatinine	Χ						X		×		×		X	
Bil (dir.)	Х						×						×	
Bii (dir.)	X													
Albumin	Χ													
PT, INR	Χ													
ALP	Х													
GGI	Х													
Giucose (f.)	Х													
HBsAg, anti-HBc (tot)	Х													
Ultrasound	X													
Fibroscan	If needed													

Where are we now?

- From 1st march, 2018 started inclusion of new patients.
- No preliminary data available for today.
- First results of project will be available for TAG-2018.

Thanks for Attention!





The 5th National Hepatitis C Elimination Workshop

HIV, TB, and HCV at Primary Healthcare Settings Pilot Project in Samegrelo-Zemo Stvaneti

NCDC 7 March, 2018



HIV, TB, and HCV at Primary Healthcare Settings Pilot Project Activities

Objective: To improve case detection of Hepatitis C, Tuberculosis and HIV infection at the primary healthcare settings

- Pilot Project
- Samegrelo-Zemo Stvaneti Region
- Program Launching 25 November, 2017



HCV,TB,HIV in Samegrelo-Zemo Svaneti Region

HCV Prevalence – 10.9% (Anti-HCV+) 7,27% (HCV RNA+)

Population - 330,761 ≥ 18 **Population** - 213,194

he Church of Martvill

Martvili

Abasha

[Georgia: 7,7 Anti-HCV+; 5,4 HCV RNA+]

Estimated number of HCV RNA+ 15,458 (23,238 Anti-HCV+)

> TB Prevalence* - 113 (100,000 population)

[Georgia 77,9 (100,000 population)]

Number of notified cases – 373 (estimated undetected cases 75(20%))

HIV/AIDS Prevalence** – 208 (100,000 population, estimated)

[Georgia:145 (100,000 population)]

- Number of cases - 868

*Database, National Center for Tuberculosis and Lung Diseases, 2017
**Database, IDACIRC



Ushgul

Need for Change

Challenges

- Significant burden of three diseases TB, HIV, HCV
- Low early detection rate and screening coverage
- Weak linkage to care and retention on treatment
- Fragmentation of services and lack of coordination

Opportunities

- Universal Health Care Program
- Universal coverage of all three diseases
- High political commitment to Hep C Elimination
- Available resources to support innovations

Strategies

- Integration of Services
- Create demand educate people, rise awareness, spread information on services
- Supply side interventions strengthening capacity of PHC service providers, community organizations to respond to challenges
- Putting right incentives for better performance



Underlined Assumptions and Key Considerations




- Disease Interactions-infection with "one" makes a person more susceptible to "other"
- Transition roots, common behavioral risk factors
- Social and economic conditions poverty, unemployment, substance use, mental illness, incarceration, social stigma, level of education
- Shared services at local level the same providers, PH, PHC, treatment sites, shared capacity
- Program Collaboration and Service Integration

CDC, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention: Strategic Plan through 2020.



Strategic Framework

A people-centred approach to care WHO




«People-centred care is focused on and organized around the health needs and expectations of people and communities rather than on patients or diseases»



Strategic Framework

A people-centred model of TB care

"An efficient and integrated set of affordable, accessible and acceptable health services, provided in a supportive environment to prevent, diagnose and treat TB"




HIV differentiated service delivery models

A client-centered approach that simplifies and adapts HIV services

"A differentiated care, is an approach that simplifies and adapts HIV services to better serve the needs of people living with and at risk of acquiring HIV and reduce unnecessary burdens on the health system"



Strategic Framework



Global health sector strategy on viral hepatitis 2016-2021



São Paulo Declaration on Hepatitis World Hepatitis Summit 2017

7. Emphasize the importance of a comprehensive, integrated and multisectoral approach that includes implementation of high-impact viral hepatitis prevention and control measures as outlined in the Global health sector strategy on viral hepatitis 2016-2021;

Need to strengthen health systems, new models of people-centered service delivery, integrated collaborative approaches and synergies between prevention and control measures for : viral hepatitis, HIV, STIs,





 To Develop the sustainable local public-private partnership for effective integration of TB/HIC/HCV screening and early disease detection

 To raise awareness among all stakeholders including general public on all thee diseases in the region











 To strengthen PHC through development of integrated TB/HIC/HCV screening model with detailed protocol







- To map the service providers at the regional level and establish effective referral systems to specialized clinics
- 200 doctors and 200 nurses, local health managers
- MoUs with private service providers






Ongoing activities

- Building the providers' capacity and increase the stakeholders' engagement in service quality monitoring;
- Meeting individually with established groups and initiate active implementation of integrated screening protocol;
- Visiting sites (including village level) for supportive supervision, individual needs assessment and additional training;
- Training of 200 doctors and 200 nurses;
- Developing municipal programs to incentivize service providers;
- Testing from April, 2018







April - October, 2018



Partners

- Ministry of Labour, Health and Social Affairs
- National Family Medicine Training Center
- National Center for TB and Lung Diseases
- Infectious Diseases, AIDS and Clinical Immunology Research Center
- Local governments
- Private service providers

Thank You!





Infection Prevention and Control in Healthcare facilities

Ministry of Labour, Health and Social Affairs of Georgia



Marina Darakhvelidze Head of Health Care Department




Infections Control Regulating Documents


- "On approval of the rules for surveillance, prevention and control of nosocomial infections" Decree of the Minister of Health №38 of September, 2015
- "On Approval of Technical Rules of Disinfection and Sterilization in Medical, Public Health and Institutions of public importance" Governmental Decree №185 of April, 2015
- Technical regalement "Management of medical waste", Governmental decree №294, 16th of June, 2017
- "National strategy on antimicrobial resistance for 2017-2020"- Governmental decree №29, 11th of January, 2017
- National guideline on infection control in medical institutions (Approved by decree of the Minister of Health №148 /o of 2009)





MINISTRY OF LABOUR HEALTH AND SOCIAL AFFAIRS OF GEORGIA







Recent legislative changes

- "On approval of assessment tool for monitoring infection control system of inpatient hospitals" Decree of Minister of Health №01-9/o, 18th of January, 2016
- "On approval of first stage plan and assessment tool for monitoring infection control system of in-patient hospitals" Decree of Minister of Health Nº01-45/0, $15^{\rm th}$ of February, 2017
- "On approval of second stage plan and assessment tool for monitoring infection control system of in-patient hospitals"- Decree of Minister of Health №01-173/o, 3rd of August, 2017.
- "On Approval of the First Round Implementation Plan of Infection Control Monitoring System in Inpatient Medical Institutions in 2018"
 Decree of Minister of Health №01-54/o, 22nd of February, 2018
- Update of infection prevention and control guideline is under development



IPC legislative Requirements Administration and enforcement mechanism

- "On Approval of the Regulations on the Rules and Conditions of Issuance of Licenses for Medical Activities and Permits of Inpatient Institutions" -Decree No. 385 of the Government of Georgia of December 17, 2010.
- "On approval of the Regulations on the Rules for High Risk Outpatient Medical Activities" -Decree No. 359 of the Government of Georgia of November 22, 2010.

It is administered by the State Regulation Agency for Medical Activities by selective control of permit/license conditions. The process also provides for the imposition of a penalties, if permit/license/high risk medical activities conditions are not satisfied



IPC legislative Requirements Administration and enforcement mechanism

- "On some measures to be taken for the purpose of transition to universal healthcare"- Decree No. 36 of the Government of Georgia of February 21, 2013
- "On Approval of Rules for Evaluation of Infection Control System in Inpatient Medical Institutions within the State Program of the Universal Health Care" -Decree of Minister of Health №01-5/n, 7 of February, 2018

The subjects, participating in the program, are obliged to take into account the recommendations issued during the monitoring of the infection control system, otherwise the participant status will be stopped

Monitoring of IPC Systems Functioning in inpatient Health Care Facilities

 Since 2016 monitoring in medical institutions of the country is conducted in the mode of assessment of the functioning of the IPC system in inpatient medical institutions. Monitoring is coordinated by the Health Care Department at the Ministry and in conjunction with the NCDC. Monitoring is carried out by the method of Supporting Supervision

Monitoring area

- ✓ Inpatient Medical Facilities ≥150 beds in Tbilisi
- ✓ Inpatient Cardiac Surgical Service providers in Tbilisi
- ✓ Third level perinatal service providers
- ✓ With the permission of Reanimation of Tbilisi Facilities ≤ 40 beds
- ✓ Inpatient medical facilities in Kutaisi and Batumi



Goals of Monitoring


On each institution level the infection prevention and control system

- Evaluation of capabilities and support supervision
- Evaluation of the necessary infrastructure and supplies for functioning
- Discover strongholds and follow it further
- Increase awareness of the requirements within the national regulations
- Find disadvantages and develop recommendations, methodological assistance in place
- Identify systemic problems and plan activities to eradicate them



IPC Monitoring results


I-II Stage (Environmental Assessment Component) 66 Establishment





IPC Monitoring results


I-II Stage (component of IPC measurement assessment) 66 Establishment



Source: IPC Monitoring results, 2016-2017



Conformity indicator with the criteria for infection prevention and control measures





Healthcare settings - Gaps

- Although regulations are in place to mandate IPC in health-care facilities, these regulations are not well enforced
- Most hospitals have no internal policy on the safety and hepatitis B vaccination for high-risk groups of HCWs (only 26% of institutions in the second stage of the monitoring have satisfied these criteria fully)
- At the second stage of the monitoring, 33% of the institutions do not meet the safeguards of safe injection practice
- The problem still remains to identify, register and report nosocomial infections by institutions, more than 50% of institutions fail to fulfill this obligation



Priorities for 2018

Improving of legal and regulatory framework:

- Develop/update the healthcare workers and patients safety policy (screening, immunization, post-exposure prophylaxis), review, recognition and implementation
- Revision and modification of the acts regulating the infection control and prevention measures in a medical institution
- Acceptance and Implementation of New Guidelines for Infection Control and Prevention
- Development of educational materials on IPC-related topics for medical personnel training of medical personnel on IPC-related topics
- In 2018, will be continue **Supportive Supervision** Monitoring of inpatient medical institutions in the country



Thank you for your attention!



Infection Prevention and Control in Georgia: Current and Future Activities

Giorgi Chakhunashvili MD, PhD
National Center for Disease Control and Public Health
Georgia



National Guideline on Infection Prevention and Control (IPC)

- In 2009, the first National Guideline on "Infection prevention and Control in Healthcare Facilities" was then developed and published in 2009. Currently, it is being updated with the support of US CDC
- Guideline on "Infection prevention and Control in Dental Settings" was then developed and published



National AMR Strategy: Strengthening IPC

Activity	Anticipated Outcome	Indicator
Establishing infection control committees at hospitals and strengthening surveillance of their performance;	Effective organizational control systems for infections at inpatient health-care facilities will be ensured	Infection control committees are established at hospitals and their adequate performance is ensured
Updating documents and preparing protocols, guidelines and other information material, regulating infection prevention and control at health-care facilities and ensuring their communication		Documents, protocols, guidelines and other information material, regulating infection prevention and control at health-care facilities are in place



Previous Awareness Campaigns on IPC

Healthcare workers

- 320 nurse received trainings on IPC (short-term)
- 40 hospital-epidemiologist received trainings on IPC (short-term)
- Long-term post-graduate educational course on IPC has been developed

Non-healthcare workers

 Training of non-medical facility staff on IPCrelated topics

General population

Mass media campaigns







IPC in Dental Settings

- Guideline on "Infection prevention and Control in Dental Settings" has been developed and implemented
- Strengthening of IPC programmes in dental settings
- About 5,000 personals of dental settings received trainings on IPC







Collaboration with Sanford Guide

- Sanford Guide to Hepatitis Therapy Android app and website in Georgian language
 - Disease characteristics
 - Information on drugs
 - Prevention
 - Dosage calculation
- Sanford Guide to Antimicrobial Therapy in Georgian language (hardcopy and web-edition) will be updated in 2018







Non-traditional healthcare and other community settings – M & E indicators and targets

- Activities to improve IPC in non-healthcare settings:
 - State regulations and policies of IPC during aesthetic and cosmetic procedures are updated and published online
 - SOPs on sterilization, disinfection, and waste management available in non-medical facilities
 - Number and percentage of non-medical facility staff trained
 - Degree to which sterilization, disinfection, and waste management SOPs are followed



The following activities are planned

- Strengthening IPC programmes in hospitals
- Surveillance of the antibiotic use
- Promotion of antibiotic stewardship
- Update National IPC guideline
- Development of IPC standard operation procedures
- Participation in media campaigns on IPC-related topics (Antibiotic awareness day, TV programs, etc.)



National Center for Emerging and Zoonotic Infectious Diseases



Implementation TAG Recommendations for Infection Prevention and Control

5th National Hepatitis C Elimination Workshop Tbilisi, Georgia March 7, 2018

Meghan Lyman, M.D.
International Infection Control Program
CDC Division of Healthcare Quality Promotion

CDC International Infection Control Program



Provide rapid assistance for outbreaks and other adverse events related to healthcare delivery




Reduce the global burden of antimicrobial resistance associated with healthcare delivery



Improve country Infection Prevention & Control capacity to prevent and control healthcare-associated infectious disease outbreaks and device-associated HAIs



Core Components of IPC



TAG IPC Recommendations

implementation of IPC all over the country.
 2. Decrees are needed to approve the country IPC structure (National, facility and primary health care levels) with clear roles and responsibilities.
 3. The central IPC authority should develop a national IPC strategy and plan to achieve implementation of IPC standards at all healthcare facilities by December 2020 (including detailed activities and timeline for implementation).

IPC Guidelines

IPC Training

& Education

 ${\bf 4. \ \ Completion \ of \ the \ } {\bf national \ IPC \ guidelines} \ {\bf and \ } {\bf making \ them \ } {\bf accessible \ to \ all \ healthcare \ providers}.$

Create multidisciplinary IPC central authority or structure (IPC unit or department) accountable for


- 5. Develop a standardized **national IPC training program** using the national IPC guidelines as the basis.
- 6. Develop a **training strategy** with the aim of targeting all healthcare providers (clinicians, nurses, others) by December 2020.
- 7. Develop and implement a **communication strategy** to promote behavior change of healthcare providers and non-traditional healthcare providers to ensure safety of practices
- IPC Monitoring & Evaluation
- 8. **Monitor implementation of IPC** at facility level by auditing IPC practices using available IPC tools to monitor progress of implementation of IPC.

IPC Program

- Progress
 - Leadership support for strengthening IPC
 - National IPC Focal Person identified
 - National action plan under development
- Challenges
 - Decentralized national IPC structure
 - Limited IPC enforcement and accountability
 - Limited resources allocated for IPC

IPC Guidelines

- Progress
 - National IPC Guidelines being developed
 - Guidelines on priority topics already approved
 - Standard and transmission-based precautions
 - Environmental cleaning
 - Equipment decontamination and sterilization
- Challenges
 - Roll-out and implementation



IPC Training

- Progress
 - Motivated IPC focal persons
 - Trained IPC champions
- Challenges
 - Limited IPC expertise
 - No IPC training strategy



IPC Monitoring & Evaluation (M&E)

- Progress
 - IPC assessments (MoLHSA)
 - Situational analysis (MoLHSA, CDC, and partners)
- Challenges
 - Lack of M&E strategy
 - Evaluating IPC implementation
 - 'Punitive' perception



Core Components of IPC

Important next steps

- Establish a strong national IPC unit with sufficient resources and authority
- Mandate IPC activities at the facility level
- Effectively roll out and implement guidelines
- Provide sufficient training to all healthcare workers
- Monitor IPC practices to guide improvement

Thank you

For more information, contact CDC 1-800-CDC-INFO (232-4636)

TTY: 1-888-232-6348 www.cdc.gov

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.




WHO GUIDANCE ON HEPATITIS SURVEILLANCE

Antons Mozalevskis, WHO Regional Office for Europe Yvan Hutin, Global Hepatitis Programme



THE GLOBAL HEALTH SECTOR STRATEGY ON VIRAL HEPATITIS



5 STRATEGIC DIRECTIONS

- 1. Information for focused action
- 2. Interventions for impact
- 3. Delivering for equity
- 4. Financing for sustainability
- 5. Innovation for acceleration

7th

leading cause of death globally (2013)



FOUR MAIN HEPATITIS VIRUSES

A. E.

В.

C.

95% of burden

Faecal oral route

Exposure to blood / body fluids

Acute hepatitis

Chronic infections

Sequelae





DATA SYSTEMS NEEDED

A | HEPATITIS SURVEILLANCE

- 1. Acute hepatitis that reflect new infections
- 2. Chronic infections
- 3. Sequelae

B|PROGRAMME DATA

- Prevention indicators
- Patient registries for the cascade of care and cure



CHALLENGES IN VIRAL HEPATITIS SURVEILLANCE

- Multiple disease outcomes
 - Consider acute hepatitis, chronic infection and sequelae
- Similar clinical presentation
 - Use in-vitro diagnostics
- Asymptomatic nature of most infections
 - Plan biomarker surveys
- Multiple modes of transmission / persons at risk
 - Tailor response to national situation



TERMINOLOGY

DISEASE

- Acute
- Chronic

INFECTION

- New
- Old
- Resolved

SEQUELAE

- Cirrhosis
- Hepatocellular carcinoma



DIFFERENTIATING INCIDENCE AND PREVALENCE

INCIDENCE

- New infections among susceptible persons
- May lead to acute hepatitis
 - Known referent exposure period (incubation)
 - Investigation of risk factors
 - Direct prevention activities

PREVALENCE

- Current infections, usually acquired in the past
- Usually asymptomatic
 - Timing of exposure unknown:
 - Investigation of risk factors limited
 - Possible to examine factors associated with prevalent infection
 - Plan testing and treatment



VIRAL HEPATITIS SURVEILLANCE

1. Detect outbreaks, monitor trends in incidence and identify risk factors for new, incident infections **2.** Estimate the prevalence of chronic infections and monitor trends in sentinel groups

3. Estimate the burden of sequelae

Surveillance for acute hepatitis

Surveillance for chronic infections

Surveillance for cirrhosis and HCC

HCC: Hepatocellular carcinoma



ACUTE HEPATITIS SURVEILLANCE

1. Detect outbreaks, monitor trends in incidence and identify risk factors for new, incident infections **2.** Estimate the prevalence of chronic infections and monitor trends in sentinel groups

3. Estimate the burden of sequelae

Surveillance for acute hepatitis

Surveillance for chronic infections

Surveillance for cirrhosis and HCC





ACUTE HEPATITIS SURVEILLANCE

	Syndromic surveillance	Enhanced case reporting
Scale	Nationwide	Mostly sentinel
Implementer	Communicable disease surveillance	Centres of excellence
Case definitions	Clinical – No in vitro diagnosis	Type specific – IgM in vitro diagnosis
Data collection	Basic demographics	Risk factors
Objectives	Detect outbreaks	Describe trends Identify risk factors



CASE DEFINITIONS FOR ACUTE HEPATITIS

Level	Acute hepatitis			
Suspect case: Clinical criteria	Discrete onset of an acute illness with signs or symptoms of (a) acute viral illness and (b) hepatic injury (liver enzymes)			
Confirmed case:	HAV	HEV	HBV	HCV
Clinical criteria AND epidemiological criteria or biomarker criteria	IgM anti-HAV + OR	lgM anti-HEV + OR	IgM anti-HBc +	Anti HCV + AND All IgM – for HAV, HEV, and anti-HBc OR
	Epidemiological link with a	Epidemiological link with a confirmed case		RNA +/ Anti-HCV – OR
	confirmed case			Sero-conversion Anti-HCV



SPECIFICITY KEY FOR ACUTE HEPATITIS

- Chronic hepatitis is more common than acute
- Only acute hepatitis can inform about risk factors
- Inclusion of cases of chronic hepatitis among cases of acute hepatitis is misleading
 - Over-estimate incidence
 - Misrepresent risk factors
- Prefer quality to quantity
 - ✓ Smaller number of sentinel sites with good diagnosis





SURVEILLANCE FOR CHRONIC INFECTIONS

1. Detect outbreaks, monitor trends in incidence and identify risk factors for new, incident infections

2. Estimate the prevalence of chronic infections and monitor trends in sentinel groups

3. Estimate the burden of sequelae

Surveillance for acute hepatitis

Surveillance for chronic infections

Surveillance for cirrhosis and HCC





SURVEILLANCE FOR CHRONIC INFECTIONS

A BIOMARKER SURVEYS

- Reference method
 - General population
 - High-risk groups

B | **DATA MINING**

Search for existing information (Gray literature)

C|REPORTING OF CHRONIC CASES

- Only estimates the quantity of cases diagnoses
- Best directed to patients registries
- Not to be mixed up with acute hepatitis surveillance



CASE DEFINITIONS FOR CHRONIC HEPATITIS

Type of criteria	HBV	НС	
Clinical criteria	Person not meeting the case definition for acute hepatitis		
Biomarker criteria HBsAg +	HBsAg +	Serological evidence of past or present infection	Chronic HCV infection
	Anti HCV +	HCV RNA + OR HCV Ag +	





04 | THREE PURPOSES – THREE DOMAINS

1. Detect outbreaks, monitor trends in incidence and identify risk factors for new, incident infections 2. Estimate the prevalence of chronic infections and monitor trends in sentinel groups

3. Estimate the burden of sequelae

Surveillance for acute hepatitis

Surveillance for chronic infections

Surveillance for cirrhosis and HCC

HCC: Hepatocellular carcinoma





ESTIMATING MORTALITY

- 1. Assess quality of the vital registration system
- 2. Group deaths directly associated with HBV or HCV infections ICD-10 codes
- Estimate cirrhosis and HCC deaths (Mortality envelope)ICD-10 codes
- 4. Estimate HBV/HCV attributable fraction of HCC and cirrhosis:
 - Centre of excellence
 - Prevalence of HBV and HCV infections in sequelae patients
- **5. Apply** attributable fractions to HCC and cirrhosis envelope
- **6.** Compile data and break down by acute / chronic and by virus



USING SURVEILLANCE TO EVALUATE PROGRAMMES

Interventions	Domain
Vaccination	Surveillance for acute
Food water safety	hepatitis that reflect
Blood safety	new infections
Condoms	
Harm reduction	
Infection control	
Testing and treating	Surveillance for chronic, prevalent infections
Hepatocellular carcinoma screening	Surveillance for sequelae
	Vaccination Food water safety Blood safety Condoms Harm reduction Infection control Testing and treating Hepatocellular

Hepatitis Surveillance Workshop

6 March 2018 Tbilisi, Georgia



What questions need to answered/addressed for HCV elimination?

- 1. Monitor the trends;
- 2. Identify the risk factors for incident infections;
- 3. Monitor the cascade of care;
- 4. Evaluate the impact of interventions.



Discussion/Considerations

• EIDSS:

- Centralized sysetm for reporting cases of notifiable conditions, including hepatitis A and hepatitis B cases
- Currenty does not collect acute HCV cases

Harm Reduction

- Incidence study /PWID cohort
- Linking harm reduction data base through unique ID with the treatment database planned

Blood Program

- Possibility for surveillance of incident infections
- Query to reveal serocenversion in the screening D/B



Discussion/Considerations

Dialysis

 HD units screen the patients, however are not mandated to report to the screening data base

Pediatric and ANC

- 2016-2017, 88,689 pregnant women tested, 437 HCV+
 (0.5%)
- From 2015, 103,399 children under 18 tested, 322 –
 HCV+
- Of 107 HCV+ childeren below 2, 54 children's mothers (50%) tested HCV- (diagnostics quality??)



Recommendations

- Strengthen human resources for hepatitis surveillance
- Establish acute hepatitis C surveillance:
 - Integrate reporting of acute HCV to EIDSS?
 - Establish sentinel surveillance in 3 ID hospitals?
- Conduct retrospectively review of ID hospitals records of acute hepatitis cases
- Link harm reduction data base with the treatment databased through unique ID, ensuring personal data protection



Recommendations

- Conduct case investigation for each HCV+ child and repeat testing at 18 months
- Ensure quality of screening tests
 - Enforce limited list of quality assured test-kits
 - Centralized provision of tests-kits
- Modify the decree to mandate HD units to report and enter all cases tested HCV+
- Use eHealth database, cancer registry and vital statistics for sequelae surveillance



Overview and Progress of the HCV Elimination Program in Georgia

Amiran Gamkrelidze, MD, PhD, Professor

Director General

National Center for Disease Control and Public Health

March 7, 2018




Nationwide HCV prevalence


- ✓ Total number of interviews –6330 (90% response rate)
- ✓ Total number of blood samples 6010 (86% response rate)

Characteristic	n	%	Estimated # nationwide ≥18 years
Anti-HCV+	433	7.7%	215,000
HCV RNA+	311	5.4%	150,000

HCV Genotype Distribution




HCV Prevalence by Age and Gender





Prevalence and Estimated Number of HCV RNA+ Individuals by Regions and Cities





Main Findings

IDU and Blood Transfusion – Most common risk factors

Highest prevalence regions – Samegrelo, Achara, Imereti, Tbilisi

215,000 anti-HCV positive persons


Men between 30-50 years old – Highest prevalence

Large proportion of individuals – unknown risk factors

Only about 1/3 of anti-HCV positive persons knew their HCV status



Road towards HCV Elimination





Road towards HCV Elimination (2)



Why Georgia?

- ► Small size of the country (69,700 km²) with population of **3.7 million** people
- ► High prevalence of HCV infection in general population
- ► Strong Governmental commitment towards elimination of HCV
- ► Availability of all modern HCV diagnostic and treatment methods
- ▶ Strong human resource capacity in the field of viral hepatitis, and particularly in hepatitis C
- ► Adherence to principles of evidence-based medicine for hepatitis C
- Existence of effective systems for implementing large-scale national and international health programs, including through multi-sectoral approach
- Availability of logistic and control mechanisms within existing national HIV/AIDS, Tuberculosis and hepatitis C treatment programs



Public/private partnership

MoHLSA

NCDC

CDC

- South Caucus Office (DGHP)
- Division of Healthcare Quality Promotion (DHQP)
- Division of Cancer Prevention & Control (DCPC)

Gilead Sciences

Others

- **Abbott Diagnostics**
- Becton Dickinson (BD)
- University of Bristol, UK
- Extension for Community Healthcare Outcomes (ECHO)
- Liver Institute & Foundation for Education and Research (LIFER)
- Global Fund for AIDS, TB & Malaria

NOhep

HCV treatment provider clinics

- Division of Global HIV & TB (DGHT)
- Division of Health Informatics and Surveillance (DHIS)
- Division of Viral Hepatitis

WHO-HQ, WHO-EURO

Foundation for New Innovative Diagnostics (FIND)

Blood Systems Research Inst.

CDC Foundation

Training Programs in Epidemiology and Public Health

Interventions Network (TEPHINET)

ENABL-ED



Goal



Elimination of HCV by ensuring prevention, diagnostics and treatment of the disease

Targets

90-95-95

By 2020

✓ 90% of people living with HCV are diagnosed

√ 95% of those diagnosed are treated

√ 95% of those treated are cured



Strategic plan for the elimination of HCV Recent progress

Promote advocacy, awareness and education, and partnerships for HCV associated resource mobilization

- Massive awareness raising campaigns
- •Short text messages (SMS) regularly sent to the population to invite them to the HCV screening
- •HCV Hotline established at NCDC

Prevent HCV transmission

- •Infection control and prevention monitoring and evaluation in medical and non-medical facilities
- •Regular screening activities among PWID and their sexual partners
- •Improved quality control mechanisms for blood banks

Identify Persons Infected with HCV

- Enhancement of screening through extending it to the more than **600 centers** countrywide, including inpatient and outpatient settings, pharmacies, Public Health Centers, Harm Reduction Cites, etc.
- Development of unified electronic screening module that combines screening data from all sources

Improve HCV Laboratory Diagnostics

- •Research activities for the simplification of HCV diagnostics, including studies on HCV cAg and various PCR approaches
- •Introduction of external quality assessment in NCDC referral laboratory (Lugar Center)

Provide HCV Care and Treatment


- •32 medical facilities providing treatment services countrywide
- •Overall SVR rate 98%

Improve HCV Surveillance

- •Identification and Characterization of HCV-Attributable Hepatocellular Carcinoma in collaboration with US CDC
- •Survey to find the barriers and facilitators to enroll in the HCV treatment program
- •HCV seroprevalence survey to be repeated in 2019




Sources of HCV screening data






HCV screening providers




HCV Screening Among Different Target Groups



Since 2015, more than 1,5 million persons have been screened

among which ~974,000 were unique individuals with positivity rate 10.6%

Screening sessions in different population groups and positivity rates



Unified Electronic Screening Module

- Real time monitoring of screening progress and data linkage with other databases
- Simplified process of getting medical services
- Easy data exchange capabilities for validation of different results
- Data standardization for better analysis
- Proper and effective validation mechanisms to avoid mistakes in business rules and data entry
- Notification and reminder systems
- Robust reporting capabilities
- Comprehensive data for improved policy making process




Blood Safety 2017 progress

- Since July 2017, participation in the Unified Electronic Donor Database became mandatory for all blood banks irrespective of their involvement in the State Safe Blood Program and for all hospitals performing blood transfusion
- Confirmatory Core-Ag testing by the Lugar Center became mandatory for all HCV screening positive blood donations
- Referral of hepatitis C screening-positive samples to the Lugar Center for confirmatory testing became mandatory for all blood establishments
- Modernization of the Donor Database has been finalized through the addition of an advanced modules to the existing core variables and linkage with the hepatitis C screening modules has been established
- An additional external quality control mechanism has been added to the existing quality control system through the professional testing by the Lugar Center
- Blood banks not involved in the State Safe Blood program were given the opportunity to participate in the program's component providing external quality control mechanism
- For the first time, mandatory voluntary donations and mandatory nucleic acid testing was introduced for blood donors with high behavioral risk




Number of HCV testing sessions among PWIDs by years






Treatment providers






Enrollment in the HCV treatment program






Georgia Hepatitis C Elimination Program Care Cascade April 28, 2015 – February 28, 2018



^{*} Of 37,500 patients eligible for SVR assessment, 28,806 were tested, 28,283 (98.2%) achieved SVR, 8,694 (23.2%) missing data




System Future Architecture






Dashboard Example from Hepatitis C Analytical Layer





Recent update in the HCV Elimination State Program

- HCV active infection confirmation testing becomes free-of-charge
- Screening providers, including inpatient clinics, blood banks, NCDC labs and public health centers will prepare venous blood samples after HCV screening positive result. NCDC will organize the transportation of the blood samples to the Lugar Center
- NCDC/Public Health Center epidemiologists commit to find and follow-up all persons with probable HCV infection





Integration of HCV, TB and HIV Detection at PHC level

«People-centred care is focused on and organized around the health needs and expectations of people and communities rather than on patients or diseases»



Pilot project in Samegrelo-Zemo Svaneti Region was lunched in September, 2017

- To raise awareness among all stakeholders including general public on all thee diseases in the region;
- To Develop the sustainable local public-private partnership for effective integration of TB/HIC/HCV screening and early disease detection;
- To strengthen PHC through development of integrated TB/HIC/HCV screening model with detailed protocol;
- To map the service providers at the regional level and establish effective referral systems to specialized clinics;
- To build the providers' capacity and increase the stakeholders' engagement in service quality monitoring;
- To generate the evidence-based recommendations for program expansion to promote service integration nationwide

HCV Elimination Program Management Ministry of Labour, Health and Social Affairs of Georgia

State Committee

Clinical Committee

Scientific Committee

National Center for Disease Control and Public Health

Screening providers - 570

Healthcare providers - 32

Social Service Agency

Hepatitis

Managament Center

- Tbilisi

Hepatitis

Managament Center
Zugdidi




Georgia HCV Elimination Clinical Committee


The American Association for the Study of Liver Diseases and the Infectious Diseases Society of America Present

HCV Guidance:
Recommendations for Testing, Managing, and Treating Hepatitis C

Lent Updated: April 12, 2017 www.bcvguidulines.arg










Georgia HCV Elimination Scientific Committee

- Established in August, 2016
- Membership
 - Ministry of Labour, Health and Social Affairs
 - NCDC,
 - Infectious Diseases, AIDS and Clinical Immunology Research Center,
 - · Clinic "Hepa",
 - Clinic "Neolab",
 - Clinic "Mrcheveli"
 - US CDC
- Open to non-members (e.g.: NGOs, Universities, other)
 - Requires member sponsorship
 - Bristol U, FIND, Boston U
- Received 40 proposals from Aug 2016 to Feb 2018
 - 40 reviewed, 32 approved



Project ECHO to Regional Facilities

(About 20 TeleECHO clinic sessions conducted through 4 hubs, June 2016 – June 30, 2017)

TBILISI

ECHO Hubs

Infectious Diseases and AIDS Center; Clinic HEPA; Clinic NeoLab; Clinic Mrcheveli

Tbilisi
3 sites

Batumi 1 site Kutaisi 3 sites Zugdidi 1 site Rustavi 1 site

Gori

1 site

Expert Panel:

Tengiz Tsertsvadze (Head of Clinical Group)

Vakhtang Kerashvili (ID Center), Elza Vashakidze (Tbilisi State Medical University), Lali Sharvadze (Hepa), Maia Butsashvili (NeoLab), George Kamkamidze (NeoLab),

David Metreveli (Mrcheveli), Maia Zhamutashvili (Mrcheveli), Jaba Zarkua (Mrcheveli), Mamuka Zakalashvili (Mrcheveli)



Important events

- National Hepatitis C Elimination Workshops 2014, 2015, 2016, 2017, 2018
- Technical Advisory Group meetings –2015, 2016, 2017





EASL side meetings dedicated to Georgian HCV Elimination
 Program

2014 – London

2015 - Vienna

2016 - Barcelona

2017 - Amsterdam

2018 - Paris



Study tours

Study tours organized by NCDC with a focus on HCV elimination.

- Introduction to the State HCV Elimination program, its management and organization
- Discussion about the steps taken to achieve the progress towards the elimination goals



Delegation from Egypt



Delegation from Post-Soviet countries

Meetings with all stakeholders:

- Ministry of Labour, Health and Social Affairs
- NCDC
- HCV Management Centers
- HCV screening and treatment provider clinics
- NCDC Referral laboratory (Lugar Center)
- Public Health Centers



NOhep Visionaries Programme



Georgia was awarded the title of NOhep Visionary for the European Region at the World Hepatitis Summit in Sao Paulo, Brazil, on November 1st, 2017.

- Six countries were selected to launch the NOhep Visionaries Programme worldwide:
 - Brazil,
 - Bangladesh,
 - Egypt,
 - Gambia
 - Mongolia, and
 - Georgia



The NOhep Visionaries Programme is a global campaign which engages governments to scale-up successful approaches to elimination and share key learnings, accelerating progress towards eliminating viral hepatitis by 2030.



Acknowledgements

















Ministry of Labour, Health and Social Affairs

US Centers for Disease Control and Prevention (CDC)

Gilead Sciences, Inc.

WHO, WHO Euro

TAG Members

LIFER

ECHO

World Hepatitis Alliance

Nohep Movement

National Center for Disease Control and Public Health



Thank you!







Management of HEPC Decentralization

Ekaterine Adamia

Head of Health Care Programs Division March 2018

Goal

Goal Elimination of HCV by ensuring prevention, diagnostics and treatment of the disease


Targets

90-95-95

By 2020

- ❖90% of people living with HCV are diagnosed
- 95% of those diagnosed are treated
- ❖95% of those treated are cured

Georgia Hepatitis C Elimination Program Care Cascade April 28, 2015 – February 28, 2018




^{*} of 37,500 patients eligible for SVR assessment, 28,806 were tested, 28,283 (98.2%) achieved SVR, 8,694 (23.2%) missing data

Challenges

- Declined enrollment in the treatment program
- Limited provider capacity and a scarcity of treatment centers in some geographic and especially in rural areas
 - Increase from 4 sites (launch of the program in April 2015) to 31 sites by October, 2017

Service provider



Objectives

- Increase geographical accessibility
- decentralization of screening, care and treatment services


- Increase financial accessibility
- > simplified diagnostics algorithms
- **➤** Minimize cost of diagnostic

Diagnostic algorithm First confirmatory test


Before 1.12.17




After 1.12.17




Diagnostic before treatment




Diagnostic for 12 week treatment monitoring



Diagnostic cost for patient Diagnostic for 12 week treatment monitoring




Diagnostic cost Diagnostic for 24 week treatment monitoring



Diagnostic cost for patient

Diagnostic for 24 week treatment monitoring



Diagnostic cost

SVR test (HCV RNA + Physician consultation) = 130 GEL

Fully Funded by Government

Decentralization First Steps

 existing regulations - all in-patient facilities are conducting HCV screening of hospital patients.

After 10 of march, they will have an obligation to conduct confirmatory testing to all HCV positive patients.


Decentralization First Steps

- The regional laboratories of the NCDC;
- The regional units of the NCDC;
- Public Health Centers;
- Harm reduction sites;




are obliged to conduct confirmatory testing of HCV positive patients;

Blood Samples Transportation



Information Flow

SCREENING MODULE!



Patient Surveillance

 From 1 of march 2018, The relevant units of the NCDC and Public Health Centers are obliged to ensure surveillance of HCV positive patients revealed by confirmatory testing and other needed diagnostic tests

Pilot Projects (1)

 "Integrating HCV screening and simplified treatment services in primary healthcare" – project of the Infectious Diseases, AIDS and Clinical Immunology Research Center, which aims to eliminate barriers and improve engagement in the entire continuum of HCV care;

Pilot Projects (2)

 "Simplifing pre-treatment diagnostic tests and treatment monitoring process in the Hepatitic C Elimination Project" - Project of Clinics "Mrcheveli" and "NeoLab"

Pilot Projects (3)

 FIND's Head Start Project, which aims to scale up effective diagnosis and linkage to care of PWIDs through harm reduction sites.

Pilot Projects (4)

Integration of HCV, TB and HIV Detection at PHC level

- ✓ To Develop the sustainable local public-private partnership for effective integration of TB/HIC/HCV screening and early disease detection;
- ✓ To strengthen PHC through development of integrated TB/HIC/HCV screening model with detailed protocol;
- ✓ To map the service providers at the regional level and establish effective referral systems to specialized clinics;

Main principles/tasks of decentralization


Integrating HCV screening and simplified treatment services in:

- Primary healthcare centers (minimum one on each municipalities)
- Harm reduction centers (minimum one in Tbilisi, one in Batumi and one in Zugdidi)
- Hospitals


Main principles/tasks of decentralization

- Simplified Treatment and Diagnostics guidelines/algorithms
- Develop reasonable minimum criteria required for facilities to provide services to HCV patients
- Initial and Continued Training for PHC, Hospital staff
- Develop necessary information systems or modules to support decentralization projects (harm reduction sites, PHC etc.) into Elim-C data base

Decentralization Project (Regions)




Decentralization Project (Tbilisi)





Patient referral system



Thank you for your attention!

Implementing HCV treatment in harm reduction centers in Georgia

Health Research Union

Maia Butsashvili



Background

- > HCV treatment was integrated with OST services at one center in Tbilisi
- ➤ Enrollment of other HR sites is necessary to treat more PWID and improve linkage to care in this high-risk group
- ➤ Technical support is necessary to facilitate the process of application to MOH for being enrolled in HCV elimination program and provide HCV diagnostic and treatment services for their beneficiaries.



Background

- From January 2018, FIND will launch the project "Feasibility, acceptability, effectiveness and cost-effectiveness of models of HCV viremia testing for confirmation and cure among people who inject drugs in Georgia"
- In arm one of this study, several HR centers will start onsite HCV RNA testing using GeneXpert platform to improve linkage to care among PWID



Pilot project

Integration of HCV treatment with harm reduction services

Funded by CDC



Goal and Objectives

The goal of this project is to integrate HCV treatment with harm reduction services at four OST/NSP sites in different regions of Georgia

Objective 1

To provide three NSP sites with security equipment to meet the MOH requirements to the facility for enrollment in HCV treatment

Objective 2

To train personnel of NSP/OST sites enrolled in HCV treatment

Objective 3

To introduce simplified testing algorithm among patients treated at harm reduction sites

Objective 4

To monitor diagnostic and treatment process and to establish ECHO teleconsultations with the Hub



Activities

Facilitate enrollment in treatment program for 4 sites with different services:

- 2 sites in western Georgia (in Zugdidi and Batumi) having on-site RNA testing (FIND study site, arm 1)
- > 1 site in Tbilisi (FIND study site, arm 1)
- One private OST site (using suboxone and not being FIND study site) with low linkage to HCV care rate



Activities

- 3 NSP sites will be provided with security equipment to meet the MOH eligibility requirements
- 1 private OST center is already equipped with security equipment
- Trainings at selected 4 sites will be conducted



Activities

- We plan to train physicians, nurses and social workers before the sites start treatment
- Each site will become the spoke of the ECHO system and will be connected to one of ECHO hubs to provide continuous teleconsultations for physicians



Simplified regulation

- This project needs simplified regulations from MOH to make possible treatment integration at HR centers
- The basic security measures (camera, security system, lockers) is necessary
- Physician on site (contracted)
- Simplified diagnostic and monitoring algorithm
- Contract with elimination program provider clinic







FIND HEAD-Start project in Georgia

François Lamoury
Senior Laboratory Expert
FIND

The 5th National Hepatitis C Elimination Workshop Tbilisi, Georgia NCDC, 7-9th March 2018





Outline

FIND HEAD-Start project

FIND activities in Georgia

- ✓ Validating new technologies for HCV diagnosis
- ✓ Pilot project with different testing pathways
- ✓ Study nested within the pilot project:

Evaluation of HCV viremia testing approaches among PWID in Georgia

✓ Cost analysis of testing pathways






Hepatitis c Elimination through Access to Diagnostics (HEAD) - Start project

Goal:

Support and build efficient and long-lasting public health response to HCV by driving policy change and accelerate access



Objectives:

- Accelerate the development of new HCV diagnostics
- Establish innovative approaches for HCV testing [screening and diagnosis]
- Assess feasibility and effectiveness of HCV service delivery approaches
- Provide evidence to scale-up differentiated HCV diagnostic services
- Contribute to global normative guidelines & national policies to support a global HCV elimination pathway





HEAD-Start: Country projects

Total Dx

Confirmatory testing: ≈36,000

Rx Targets

Treatment: ≈7,200 (FIND), ≈20,500 (Partners)

Georgia:

Provide confirmatory testing for 10,500

Myanmar

Vietnam

Malaysia

Incidence of HCV infection

5.6-6.6

1751

21.3-28.7

Total number (000)

104-124

1572-2120

Cameroon

India: Delhi Manipur Punjab WHO region Uncertainty Map key Uncertainty estimate interval African Region 31.0 22.5-54.4 222-544 Region of the Americas 6.4 5.9-7.0 59-69 Eastern Mediterranean Region. 62.5 55.6-65.2 365-426 European Region 61.8 50.3-66.0 460-603 12.5-26.9 South-East Asia Region 14.8 243-524

6.0

23.7

Incidence rate (per 100 000)

Source: WHO, work conducted by the Center for Disease Analysis. See Arress 2.

Western Pacific Region

These deaths exclude these from the antishapatic complications of HCV information and background mortality from other passes arrons HCV. Informational configuration.





Expanding new technologies for HCV testing Validation project - Georgia

Xpert HCV VL FS assay

Easier blood collection by fingerstick

Result in one hour







Study closing @ Lugar Center & Hepa+, recruited 400 participants.

Other countries: Cameroon, Malaysia, Greece, Denmark

Future potential projects:

Use of Dried Blot Spots for HCV RNA testing

Rapid Diagnostic Test evaluation for more quality assured products





Pilot project - Georgia



Increase linkage between screening and confirmatory testing

Help move elimination of HCV in Georgia forwards






FIND pilot project- scope

ACTIVITY

OUTCOME



Decentralization of HCV confirmatory testing at HRS using **GeneXpert plasma** cartridges





 Support the integration of HCV RNA testing in TB GeneXpert

Decentralization of HCV confirmatory testing at HRS using **Abbott cAg assay** at Lugar Centre



Effective sample referral mechanism and efficient pts workflow increases access to HCV confirmatory and decreases TAT




 One testing approach in high prevalence setting (modelling)





Overview FINDs contribution







STUDY - portion of pilot project

Evaluation of HCV viremia testing approaches among PWID in Georgia

Goals

 Service delivery approaches to provide confirmatory testing for a greatest number of PWID being:

Screened diagnosed Completing treatment

- Use of cAg for test of cure (SVR 12)
- Cost of the service delivery approaches; core antigen and point of care RNA





HRS sites included in the pilot project





Point of Care at HRS



cAg, samples collected at HRS and transported to Lugar Center



Standard of Care, patient referred for dx and tx



Study methods

Harm Reduction Sites

A A A A O D D D

Non randomized assignment

Arm 1 - POC



620 beneficiaries who screen HCV+

- Blood draw
- HCV RNA testing
- Results to PWID

Arm 2 - cAg



620 beneficiaries who screen HCV+

- Blood draw
- Blood sample shipped to Lugar
- HCV cAg test
- Results to PWID cAg as test of cure

Costing of pathway

LTFU rate along care cascade

Arm 3 - SOC



620 beneficiaries who screen HCV+

- Current practice





Substudy: Evaluation of the diagnostic performance of HCVcAg as test of cure in for hepatitis C among PWID in Georgia

- HCVcAg: highly sensitive and specific in the diagnosis of active HCV infection in different populations.
- HCVcAg confirmatory testing already approved by the Georgian government
- **Limited data** on its clinical utility to confirm sustained virologic response (**SVR**) in people who have completed therapy for hepatitis C.
- Hypothesis: The use of HCV cAg assay may present an alternative to confirming SVR.
- 1^{ry} objective: To estimate the diagnostic accuracy of HCVcAg assay in differentiating patients who have achieved sustained virological response (SVR) from those who fail SVR among those who have completed treatment for hepatitis C.
- Analysis plan: The performance of the HCV cAg Assay to confirm SVR will be evaluated in comparison with Abbott *RealTime* HCV viral load assay HCV. (sensitivity, specificity, PPV, NPV. Subgroup analyses will include performance of HCV cAg by: HIV status, HBV status, HCV genotype)




Cost analysis of testing pathway






Arm 2 - cAg



Arm 3 - SOC




L1: primary health care

L2: Secondary health care L3: Tertiary Health Care





Georgia pilot project timelines





THANK YOU!

Special thanks to our partners in this endeavor!



















We are grateful for the input and feedback of many of the organizations also doing great work in the area of HCV elimination in Georgia












Study of Barriers for HCV treatment involvement in Harm reduction Centers

Maka Gogia


Georgian Harm Reduction Network March 9, 2018



Needle and Syringe Program's Geography in Georgia



Number of HCV testing sessions among PWIDs by years





Harm Reduction program data, 2017

City	Harm Reduction Site	HCV Screening	HCV positive cases	No RNA diagnose	RNA tested and involved in treatment	
	New Vector	1402	516	322	60%	
Tbilisi	New Way	2284	661	425	65%	
	Нера+	2393	672	135	35%	
	Aceso	2132	576	282	51%	
Kutaisi	New Way	2683	1193	955	20%	
Zugdidi	Qsenoni	1795	708	212	90%	
Batumi	Imedi	1414	467	210	35%	
Telavi	Step to Future	1015	240	141	38%	
Gori	Step to Future	1645	499	272	55%	
Poti	Ordu	751	277	168	43%	
Ozurgeti	Ozurgeti Fenix		242	20	90%	
Samtredia	Samtredia New Way		245	200	47%	
Rustavi	New Vector	1935	513	178	50%	

To address the barriers within HCV Elimination program

- In 2016 GHRN conducted qualitative study to assess First Phase of HCV Elimination program
- In 2018 GHRN together with partners plans to carry out quantitative study

Title of project:

Study of barriers to enrollment into HCV elimination program among PWID

Part of NCDC Study of barriers among general population

Objective of the proposed research is to identify barriers of enrollment into HCV elimination program among HCV antibody positive people

Duration:

9 months (April-December, 2017)



Collaborators and responsibilities

Organization	Responsibilities	
Health Research Union	The project lead (PI) - plans and supervises all project activities, participates in developing research design/protocol, data analysis, development of recommendations and preparation of research papers	
Georgian Harm Reduction Network	Ensures Interviewers' selection and trainings; Conducts on-site monitoring visits at 13 harm reduction sites; ensures high quality of quantitative data	
Medecins du Monde, France	Support in development of study design, data analysis and manuscript preparation	Funders
US CDC	analysis and manascript proparation	
NCDC		



Methods

- The data that are entered into the elimination program databases will allow tracking each individual from screening to the sustained viral response (SVR) test.
- The stratified sample of PWID having positive HCV antibody test between April-September, 2017 and not linked to care will be selected.
- Individuals who are lost to follow-up are defined as those not having HCV RNA or Core antigen test in the treatment database at 90th day from being screened HCV positive.
- A "linked to care" individual is defined as: HCV+ screening and having RNA/Core antigen test in the treatment database.



Methods

- Proportional to size sample of lost to follow up as well as linked to care PWID from each HR site will be selected.
- Overall sample size will be 500 PWID.
- The telephone interviews will be used.
- Interviewers will be recruited from HR center staff (HIV voluntary counseling and testing counselors) and trained prior to the start of the study.



Methods

- Five attempts to reach each study participant will be made, as done by NCDC barriers' study interviewers.
- Failure to reach the respondent as well as refusal rate will be calculated.
- The questionnaire will be piloted and adapted accordingly.
- The questionnaire will be similar to the one used by NCDC, with few additional questions specific to PWID




Expected outcomes

 We expect the study results will generate important data to understand barriers of linkage to HCV care among PWID.

 This will make possible to plan effective interventions to improve the enrollment into HCV elimination program ensuring better coverage of most important and hard to reach population as PWID to achieve HCV elimination goal.






C Elimination

Thank you!

Progress Made in Assuring Quality HCV Diagnostics in Georgia

BETH SKAGGS, PHD
CDC SOUTH CAUCASUS OFFICE

Access to Accurate and Timely Laboratory Diagnostics is the Backbone of the HCV Elimination Program



- ■In 2015-2016 ~600,000 HCV related tests were performed in Georgia
- ■US reports ~1-3% error rate among all clinical laboratory tests
- ■A 2% error rate of HCV screening in Georgia would result in 12,000 incorrect HCV tests over this 2 year period
- •Quality Management System is intended to detect errors, prevent them in the future and reduce cost

Where We Were..... *Challenges Presented in April 2016


- Appropriate laboratory services in regions yet to be established
 - Services in Tbilisi are adequate
- No unified external quality control system exists
 - Quality control limited to internal control procedures
- Test-kits used by laboratories vary based on availability from the suppliers; most of test-kits have not been validated
- •Unified standard procedures not identified for participating labs
- No national certification systems for clinical laboratories exists
- No systematic training of lab personnel in biosafety standard
- Screening HIV antibodies not part of program

Current Situation...March 2018


- Regulatory Basis for Laboratory Quality Assurance
 - Elimination Plan Strategy 4: Improve HCV Laboratory Diagnostic
 - Identification of National Reference Laboratory (NRL)
 - Creation of a National External Quality Assurance Program for PHL, Clinical and Blood Bank Laboratories
 - ➤ MoH decree "On provision of Lab Quality Assurance for the Hepatitis C elimination program"
 - Minimum requirements and mandatory tests for lab providers participating in hepatitis C elimination program
 - Development of SOPs and implementation in the participating labs
 - National EQA Program Structure
 - ► Laboratory Regulation effective January 2017

Expanded Access to HCV Laboratory Diagnostics

16 HCV PCR Laboratories (14 HCV Genotype); 1 HCV Core Ag Laboratory



Establishment of a National EQA Program



Role of NRL in Establishment of National External Quality Assurance Program

- Demonstrate acceptable performance in International EQA Programs
 - HCV VL and genotyping
 - Serology for HBV, HCV, and HIV
 - HCV Core Antigen
- Maintain and expand ISO 15189 accreditation
- Develop Georgian proficiency testing (PT)Panels
 - Agreement with Blood banks
 - Characterization, prep and store of >600 PT aliquots
 - Monthly stability testing of PT aliquots
 - Routine monitoring and documentation of internal quality control





Role of NRL in Establishment of National External Quality Assurance Program

- Develop EQA Program Documents (dual language)
 - Panel Instructions for participants
 - PT Survey forms
 - Panel Receipt Confirmation forms
 - Individual and Summary Reports
 - Feed Back
 - Data analysis and recommendations
 - Site visits
- Continue to enroll laboratories
- Provide Result Reports to participating labs and follow up support as requested
- Retain confidentiality and secure storage of EQA
 Program Data





Perticipating Laboratory Name and Address	
VERSI	
v J.TE, Vacha-Pshanela Str., 0360, Tbilisi	
una develotze@wwmi.go	
	Securitying External Guality Forms
hank you for perticipating in the Hapathin C Viral Load &	
hank you for perticipating in the Hapathin C Viral Load &	
hank you be pericipating in the Hapathia C Viral Load & v Vograms. Listed halves are the expected results for the pro- Neur lab redies	
hank you he perioquiting in the Hapathis C Viral Load & Vograms. Listed listing are the expected marks for the pro-	002

GENATOVL 2017

No.	Specimens and Lot represent	VL samps (Log (G)*	Vous teach Detected VL GogSFR	Doubletten	Seradat
3	SENATOVI OL LOTZOS 70000	Positive 4.67-4.63	Partie 631	Acorphytile	1.4
2	GENATOVI 02	Negative \$20-500	Negative \$40	Territori.	
1	GENATICAL 00	Fortier &49-647	Fodtie 5.96	Good	. 3
+	GENATOVI DE 10720/2006	Portion 4,854.55	Politice 6.15	Good	10
3.	GENATOVLOS LOTAGORAGO	Postine \$464.68	Fortier 429	Good	

Hepatitis C Viral Load Results

Georgian National PT Panel for Viral Load and Genotyping

- Locally representative source
- Each panels includes 5 specimens
- Coding of PT specimens is unique for each laboratory to avoid consultation of obtained results between labs prior to submission
- Distributed 3 times per year
- Data from the participating laboratories analyzed for feedback
 - Individual Laboratory and Summary Reports



National EQA Program Results 2017

Expansion of Laboratories has Created a Diversity of Reagents, Kits and Equipment

	Viral Load (16 labs)	Genotyping (14 labs)
RNA Isolation Kits	9	10
PCR Assay Kits	6	7
PCR Equipment	9	8

^{*}Currently no data on types of Qualitative HCV kits in use









Methods for PT Results Analysis

- All viral load results are quantitative and reported in log₁₀
- •For each PT specimen, **Mean and Standard Deviation** (SD) of quantitative VL results are calculated from all laboratories, including Reference Laboratory

```
    +/- 1 SD - Excellent
    +/- 2 SD - Good
    +/- 3 SD - Acceptable
```

- Two laboratories do not perform Genotyping
- In December 2017 PT Round, 2 laboratories did not participate citing insufficient funding/reagents to do so

EQA Program Results 2017

Table 1: QUANTITATIVE VIRAL LOAD

PT Round	# of Labs Reporting	Total # of VL Results	Percent within 1 SD - Excellent	Percent between 1-2 SD - Good	Percent between 2-3 SD - Acceptable
July 2017	13/13	65	80%	14%	6%
Sept 2017	15/15	75	75%	19%	6%
Dec 2017*	13/15	65	75%	21%	3%

Table 2: GENOTYPE

PT Round	# of Labs reporting	Percent with correct result	Notes			
July 2017	11/13	100%	2 labs do not perform test			
Sept 2017	13/15	100%	2 labs do not perform test			
Dec 2017*	11/15	100%	2 labs do not perform test			

^{*}Two enrolled laboratories did not participate in Dec PT round citing excess costs to do³so

Specimen 01 – HCV Viral load (comparative)

Obtained results (log10) (July)

Lab	Ref	001	002	003	004	005	006	007	800	009	010	011	012	013	*	*
SD	-1 SD	+1SD	+3SD	+1SD	+1SD	+1SD	+2SD	+1SD	-2SD	+1SD	-2SD	+1SD	-2SD	+1SD		_
log10	5.34	5.69	6.31	5.70	5.72	5.79	6.00	5.88	5.30	5.67	5.06	5.58	5.29	5.48		
Result	Excel	Excel	Accept	Excel	Excel	Excel	Good	Excel	Good	Excel	Good	Excel	Good	Excel		


Obtained results (log10) (Sept)

Lab	Ref	001	002	003	004	005	006	007	008	009	010	011	012	013	014	015
SD	-1SD	-1SD	+1SD	+1SD	-1SD	+1SD	+1SD	+1SD	+2SD	+1SD	-2SD	+2SD	-2SD	+1SD	-1SD	+1SD
log10	5.27	5.47	5.84	5.74	5.22	5.76	5.74	5.76	5.87	5.69	4.80	6.07	5.03	5.62	5.43	5.65
Result	Excel	Good	Excel	Good	Good	Good	Excel	Excel	Excel							

Obtained results (log10) (Dec)

Lab	Ref	001	002	003	004	005	006	007	008	009	010	011	012	013	014	015
SD	-1SD	-1SD	+1SD		-1SD	-1SD	+1SD	+1SD	+1SD	+2SD	-2SD		-3SD	-1SD	-1SD	+2SD
log10	5.35	5.42	5.85		5.56	5.52	5.57	5.99	5.78	6.16	5.06		4.71	5.46	5.46	6.30
Result	Excel	Excel	Excel	N/A	Excel	Excel	Excel	Excel	Excel	Good	Good	N/A	Accep	Excel	Excel	Good

Cumulative 2017 EQA Program Results-Quantitative HCV Viral Load



Qualitative Interpretation of HCV Viral Load Results - 2017

PT Round	HCV RNA Qualitative interpretation	HCV Genotyping						
	# of Labs reporting Percent correct results	# of Labs reporting Percent correct results						
July	13	11						
2017	100%	100%						
Sept	15	13						
2017	100%	100%						
Dec	13	11						
2017	100%	100%						

- No false negative results
- No false positive results
- No errors with Genotype

Conclusions and Recommendations

- Significant diversity of the reagents and equipment
 - FDA-approved automated tests most accurate
- Some labs modify test methods without validation
 - All labs should follow the manufacturers recommended protocols, conduct and document test validations
- Some laboratories have switched to qualitative PCR for confirmatory testing
 - EQA Panels need to be modified to better control for qualitative PCR testing and interpretation
- HCV Rapid Testing are not quality assured
 - Centralize procurement of all HCV rapid tests and include verification of tests in NRL before confirming purchase
- Standardized package of SOPs for quality management and biosafety are not available

Where We Were..... *Challenges Presented in April 2016

- \blacksquare Appropriate laboratory services in regions yet to be established \bigvee
- ■No unified external quality control system exists
- ullet No national certification systems for clinical laboratories exists $oldsymbol{\mathsf{V}}$
- ■Screening for HIV antibodies not part of program
- Test-kits used by laboratories vary based on availability from the suppliers; most of test-kits have not been validated X
- Unified standard procedures not identified for participating labs X
- No systematic training of lab personnel in biosafety standard X

Next Steps

- Expand pool of PT specimens to include additional genotypes; and new tests: qualitative PCR and GenExpert
- Establish HCV serology EQA panels and program for blood banks
- Standardize specimen rejection criteria for referred specimens to quickly identify and correct problems in specimen collection and transport systems
- Include hematology and biochemistry EQA results in National EQA database for all labs in elimination program
- Conduct laboratory assessments, including review of test validation documentation
 - Document progress in labs assessed in 2015
 - Establish baseline data for new labs since 2015

Remaining Gaps

- •Quality Assurance System for HCV Rapid Tests
 - Funding for centralized procurement and supply chain management
- Package of standardized SOPs
 - use and documentation of IQC
 - test validation
 - Root Cause Analysis and Corrective and Preventative Action
 - management of infectious waste
- Standardized training in QMS and biosafety
 - To support compliance with laboratory certification standards

Acknowledgements

Dr Jan Drobeniuc, Division of Viral Hepatitis, CDC

Dr Radu Cojocaru, Asst Prof, School of Public Health, Moldova

Dr George Kamkamidze, Director, NeoLab

Dr Maia Alkhazashvili, Manager, Lugar Center, NCDC

Dr Roena Sukhiashvili, Mol Dept, Lugar Center, NCDC

Dr Gvantsa Chanturia, Head of Mol Dept, Lugar Center, NCDC

Dr Nazibrola Chitadze, Head of Serology Dept, Lugar Center, NCDC

Nino Berishvili, HepC EQA Program, Lugar Center

Laboratory service providers of the HCV Elimination Program



Decentralization of HCV Care: Simplified Diagnostics and Treatment Monitoring

Karla Thornton, MD, MPH
Professor, Infectious Diseases
University of New Mexico
Associate Director, Project ECHO
09 March 2018



Overview

- Importance of decentralization of care to achieve elimination goals and address the TAG recommendations
- Real world examples of decentralization and simplified algorithms
 - New Mexico, United States
 - Punjab, India

Why Decentralization? TAG Recommendations to Meet Elimination Goals

- Rapid increase in number of providers in primary care trained to test and treat HCV
- HCV Treatment sites should include all hospitals, prisons, harm reduction programs, medication assisted treatment centers and TB and HIV treatment sites
- Referral relationships should be formalized and strengthened between the current major HCV treatment sites and non-tertiary care providers integrating HCV testing and treatment services in clinical settings
- The major HCV treatment sites can provide mentors and educators for newly trained providers
- ECHO provides a platform to strengthen educational, training and referral relationships

Decentralization and Simplified Treatment Algorithms

- There is not one accepted, tested way to implement decentralization
- There is not one accepted, simplified treatment algorithm
- Some key principles
 - Expands access to care
 - Better serve individual needs (treat closer to home or in acceptable setting) – integrated service delivery, differentiated care
 - Minimize expensive diagnostics (e.g., use pangenotypic if available to avoid genotyping)
 - Minimize labs without sacrificing safety

Georgia: Projects Underway

- Simplified diagnostics and treatment monitoring of HCV infected patients in rural primary care facilities in the country of Georgia
 - Tengiz Tsertsvadze, IDACIRC
- Simplification of pretreatment diagnostic evaluation and on-treatment monitoring procedures within HCV Elimination Project
 - Jaba Zarqua, Mrcheveli

Real World Examples

- New Mexico, United States
- Punjab, India



Moving Knowledge Instead of Patients

New Mexico

- Low population density, large geographic area
 - 315194 km²
 - 2.086 million people
- Most areas have few health care providers and no specialists







Hepatitis C in New Mexico (2004)



- More than 35,000 reported HCV cases
- Less than 5% had been treated
- Only one academic health center treated HCV
- Highest rate of cirrhosis deaths in the nation

Goals of Project ECHO

- Develop capacity to safely and effectively treat HCV in all areas of New Mexico and to monitor outcomes
- Develop a model to treat complex diseases in rural locations and developing countries

Methods

- Use technology to leverage scarce healthcare resources (specialty knowledge and expertise)
- Share "best practices"- reduce variation in care
- Case based learning (learning by doing)
- Centralized database to monitor outcomes


Steps

- Train physicians, nurses, pharmacists and their teams in HCV care
- Conduct teleECHO clinics "Knowledge Network"
- Initiate case-based guided practice "Learning loops"
- Collect data and monitor outcomes centrally





Learning Loops

- ✓ Interactive Learning Environment
- ✓ Co-management of Cases
- ✓ Learning by doing
- ✓ Learning from didactics
- ✓ Learning from each other
- ✓ Collaborative Problem Solving



ECHO vs. Telemedicine











Project ECHO Spoke Participant




Project ECHO Hub



Hepatitis C Genotypes 3 Treatment Regimen Decision Tree



Genotype 3 Patients




Project							for 12 weeks	
(ECHO)					Hepatitis C N	/linimum Visit/	imum Visit/ Labs Flow Shee	
		Wk 0	Wk	Wk	Wk	Wk	Wk	
Week of Treatment	Screening	Start of Tx	2	4	8	12 End of Tx	24	
Dates	N/A	01/01/18	01/15/18	01/29/18	02/26/18	03/26/18	06/18/18	
Visit		х	x	x	х	х	х	
HCV RNA	x						x	
CBC w/ Diff	х			х	х	х	x	
Chem 7	x			x	x	x	x	
LFTs/HFP	x			x	х	х	x	
HBsAg HBs Ab HBc Ab*	х							

Key Points to Remember:

- 1) Week 0 Visit is the day of the first dose of medication.
- 2) Lab draws are done the end of the treatment week.
- HBc Ab should be total or IgG.

Patient Name:		Date of Birth:			Patient ID:	Genotype:	
							REV: 11/20/17 SLS

Hepatitis C Treatment in New Mexico



Tackling the Hepatitis C Disease Burden in Punjab

- June 2016 Government of Punjab launched a public health program under the Mukh Mantri Punjab Hepatitis-C Relief Fund (MMPHCRF) that offers free HCV treatment to all residents of Punjab
- The goal is to eliminate HCV from Punjab in 10 years
- Treatment offered through a decentralized network of 22 District Hospitals and 3 Government Medical College
- 250 clinicians have been identified for training on diagnosis and treatment of chronic HCV


Burden of Hepatitis C in Punjab



- Prevalence 3.29%
- Anti-HCV positive ~920,000
- Viremic ~600,000

Demography

- Districts 22
- District Hospitals22
- Sub Divisional Hospitals 41
- **CHCs** 150
- Govt Medical Colleges 03



Capacity Building of HCW

- Hep C Workshop 4 hour course
 - 80 Clinicians trained from 25 centers (3 GMCs
 + 22 District Hospitals)
- ECHO
 - Hub: PGIMER, Chandigarh
 - Spokes (25 centers)
 - Syllabus
 - Case discussions
 - Didactic lectures
- What's App Group

ECHO Session-Spokes




Assessment Prior to Treatment (HCV RNA +)

- Detailed history and physical examination
- Baseline tests
 - CBC, LFT, RFT (Renal Function Tests)
- Viral co-infections
 - HBsAg, HIV
- Ultrasound of the abdomen
- Women of childbearing age
 - Urine pregnancy test
- Evaluation for diagnosis of cirrhosis

Evaluation of Cirrhosis


- **Fibroscan** Cirrhosis (METAVIR F4): ≥12.5 kPa
- If Fibroscan not available:
 - **Platelets** <75,000/cmm
 - APRI ≥ 2.0
 - FIB-4 > 3.25




Follow up and Tracking of patients

- Treatment Cards for patients availing MMPHCRF
- Master Data with UID for each case: Aadhar based
- Follow up of case: Every 4 weeks in order to counsel the patient and to check the patient
- Centralized case record
- SVR (Sustained Viral Response) records: 12
 weeks after treatment completion

Cascade Report (June 2016 - Jan 2018)



Pakistan



Gower E, Estes CC, Hindman S, Razavi-Shearer K, Razavi H. Global epidemiology and genotype distribution of the hepatitis C virus. J Hepatology 2014.

Replication of the ECHO Model in Georgia

Project ECHO and the Georgia HCV Elimination Program



January 2016
 Albuquerque, NM

First TeleECHO Clinic









Georgia HCV Elimination Program



Summary

- Pioneering work
- Need for ongoing evaluation and quality improvement
- Decentralization need for intense coordination
- Need ongoing education/mentorship of providers
 - Dissemination of new information
 - Complexity of cases
 - Referral pathways

Slide Acknowledgements

• Rhada Diman, MD

Scientific Committee Progress

Tinatin Kuchuloria, MD, PhD, MSc CDC Foundation

Establishment of the Scientific Committee

- Established in August, 2016
 - Research agenda development
 - Transparency
 - Coordination
 - Communication
- The need for such a committee was to
 - Avoid overlapping efforts
 - Assist stakeholders in data analysis
 - Assist in dissemination of research findings
 - Obtain resources for conducting priority activities







Scientific Committee

- Co-chaired by NCDC and CDC representatives
- Members
 - Molhsa
 - NCDC
 - IDACIRDC
 - Clinic Hepa
 - Clinic Neolab
 - Clinic Mrcheveli
 - US CDC

Members and Non-members

- Members
 - Clinicians, researchers, policy-makers
 - Local and international stake-holders
- Non-members (e.g.: NGOs, Universities, other partners) may be invited ad hoc
 - University of Bristol
 - FIND
 - Boston University Medical Center
 - LIFER
 - ECHO
 - MDM
 - EASL Foundation

Roles of the Scientific Committee

- Review, approve, and support study proposals
 - IRB review and approvals
 - Assist in securing funding
 - Study implementation
 - Data analysis
 - Manuscript writing
- Coordination with
 - Molhsa
 - Clinical Committee
 - International organizations
- Journal club

Scientific Committee Progress (August 2016 – February 2018)

- Number of proposals reviewed/approved:
 - 40 reviewed
 - 32 approved
- Types of proposals approved
 - Clinical outcomes (treatment/SVR, long-term survival, other)
 - Linkage and access to care
 - Innovative diagnostics
 - Simplified regimens and treatment expansion
 - HCV incidence studies
 - Molecular epidemiology
 - Establishing a specimen repository
 - Modeling

Scientific Committee Progress – Funding

Proposal Type	Funded	Funding required	N/A	Total
Epi studies	5	4	2	11
Database Analysis	1	0	9	10
Diagnostic studies and sample archive	4	0	0	4
Healthcare Evaluation	5	0	0	5
Modeling & Economic Evaluation	1	1	0	2
Total	16	5	11	32

Presentations and Publications

Oral and Poster presentations

- EASL, 2017 9 poster presentations & Georgia session
- AASLD, 2017 5 poster presentations
- WHS, 2017 7 poster presentations & Georgia session
- EASL, 2018 6 poster/1 oral presentations & Georgia side-meeting

Manuscripts

- Already published 4
- In press 2
- Finalized and submitted to the peer reviewed journal 1
- Drafted 1


Screening & Linkage to Care

A Key Challenge

Improving Access to Care and Treatment

- Recently approved proposals to improve access to care and treatment:
 - Assess effectiveness of screening and linkage
 - Assess barriers
 - Decentralization of services into primary care and harm reduction settings

Documenting HCV Elimination



The Role of Research in Public Health Programs

- "The research-policy arena is assumed to be a retail store in which researchers are busy filling shelves of a shop front with a comprehensive set of all possible relevant studies that a decision-maker might some day drop by to purchase.1"
- Co-production of research knowledge increases
 - Relevance
 - Comprehensibility
 - Likelihood of use of a piece of research

¹Lomas J. Connecting research and policy. Can J Policy Res. 2000;1:140–144



Conclusions

- Coordination of efforts improves efficiency
- Robust research agenda is critical to support elimination program goals
- Lessons learned from the Georgia HCV Elimination will guide other elimination efforts globally

Acknowledgements

- Ministry of Health, Labor and Social Affairs of Georgia
- National Center for Disease Control and Public Health of Georgia
- Infectious Diseases, AIDS and Clinical Immunology Research Center, Georgia
- Clinic Neolab, Georgia
- Clinic Mrcheveli, Georgia
- Centers for Disease Control and Prevention, USA
- CDC Foundation, USA
- Non-member organizations: ECHO, University of Bristol, FIND, others

Thank you

- "If it can't be done in Georgia, it can't be done."
 - Francisco Averhoff



"Hepatitis C Cured Patient Association"



Founders:

Mr. Temur Radiani, Doctor of Biology, Author of the idea;

Mr. Irakli Koridze, an economist;

Mr. Ezekiel Intskirveli-Chkheidze, Head of Kvemo Kartli Forestry Service;

Mr. Beso Kupreishvili, Founder and Director Of Finger Theater.



Introduction



Mission

Promoting successful implementation of HCV elimination program in Georgia, Raising awareness on viral hepatitis and reduction of stigma and discrimination related to hepatitis in society.







Goals (I)

- 1. Uniting the Patients Cured from HCV and forming a supportive advocacy group;
- 2. Raising awareness on HCV elimination program in the population;
- 3. Reducing stigma and discrimination related to hepatitis in medical institutions and societies;
- 4. Attracting population to increase awareness on the problem of hepatitis C;

Goals (II)

- 5. Cooperation with governmental, non-governmental organizations and funds to support the communication activities within HCV elimination program;
- 6. Increasing popularity and availability of the successful programs and disseminating information on new initiatives; cooperation with the EU professional institutions;
- 7. Sharing the Information and Experience on HCV Elimination in Georgia Participating in international meetings and conferences organized by international advocacy organizations;
- 8. Expansion of Association by increasing the involvement of cured patients.



Suggestion's

Changes in Legislation/Mandatory screening

• Strengthening Infection Control in Health Care and Non-Medical Facilities





WORLD HEPATITIS DAY

JULY 28TH

Baby boomers

[1945]
to
1965
more likely to
have HCV than
other adults


75% of adults with HCV in US are baby boomers 325 million people with chronic hepatitis infections worldwide in 2015 80% of liver cancer cases in the world 90% of people with HBV and 80% with HCV are unaware of their

status

30 to 45 seconds a person dies from HBV

Injecting drug users account for at least to de HCY transmission in the US

Republic Efficies «H



Memorial Day ???



"In my struggle for my life, I've learned that whatever happens to you, you should never give up, you should keep fighting, and eventually with a little luck and a lot of help from kind people, the virus will be defeated! I am sure that a few more years will pass and we will all be celebrating the full elimination of hepatitis C in my country,".

Temur Radiani, cured of HCV, Georgia

Thank you for attention!

Questions?

5th HEPATITIS C

Technical Advisory Group

TAG Meeting

February 9, 2018

blood Safety

Eteri Kipiani

National Center for Disease Control and Public Health



Outline

- Status of blood safety program in blood banks
- Progress
- Challenges
- Next steps



Current Status Distribution of Blood Banks in Georgia, 2018

22 blood establishments hold state license in blood production service, 15 blood banks out of them participate in the State Safe Blood Program (2 additional banks have been involved since 2018)






Current Status Country Requirement, 2017



Blood donation by 1% of the population can meet a nation's most basic requirements for blood

Percentage of Georgian population donating blood






In 2016, a total of 53000 blood transfusions were performed by hospitals

In the same year, 1,4% of Georgian population donated blood and more than 80000 donations were collected by blood banks



Current Status Country Profile

- Blood donation rate is higher in high-income countries, while prevalence of Hepatitis C shows inversely proportional trend
- Georgia is between upper and lower middle income countries in blood donation rate and between lower middle-income and low-income countries in the prevalence of Hepatitis C






Statistical Data, 2015-2017 Voluntary vs. Paid vs. Family/Replacement

Based on analysis of statistical data from 2015-2017:

- Paid donations prevail on the number of non-remunerated and family/replacement donations
- Repeat donors constitute the majority of donations
- Total number of donations has increased by 12% during last three years with slight increase (up to 5%) in the number of non-remunerated donations






Current Status Age and Gender Distribution, 2017

More than 100 000 donations have been made at all blood banks (including those not participating in the program):

- Male donors constitute majority (58%) of the total donors
- Highest prevalence was observed among males in age group of 40-49 and 30-39






Current Status First-time vs. repeat donors among anti HCV positive donations

In 2017, a total of 78 156 donations have been performed by blood banks participating in the program, including 583 anti HCV positive donations with the highest rate in first-time family/replacement and first-time voluntary unpaid donors and the lowest in paid and voluntary unpaid repeat donors








Curt Status Anti HCV prevalence by gender and donor status, 2015-2016

According to the statistical data analysis of 2015-2017, first time replacement and voluntary unpaid male donors prevail on the number of repeat

Anti HCV Prevalence in Repeat and First-time Donor Population by Gender and Donor Status, 2015-2017






Current Status

Distribution of Anti HCV Positive Donations in the Regional Context

In 2017, highest rate of antibody positive donations were registered by blood bank located in Samegrelo region showing direct correlation with national burden of HCV spread in Georgia





Progress Amendments to State Safe Blood Program

External Quality Control System

- Blood banks that do not participate in the program were given the opportunity to participate in the external quality control system
- An additional professional testing mechanism has been added through the Lugar Center
- Two additional specialists will be assigned to draw out SOPs and provide monitoring and inspection of blood banks

Confirmatory

Testing

Blood banks must ensure:

- Transportation of Hepatitis C screening-positive samples to the Center for confirmatory testing using Cor-Ag method
- Transportation of HIV screening-positive blood samples to AIDS Center for confirmatory testing
- Mandatory notification of screening positive blood donors about their status
- Removal of blood and blood components from previous donations of screening positive donors

Financing

- All participating blood banks will receive 17 GEL for each donation
- Quarterly, the blood banks will receive additional 2 GEL for each donation if proportion of voluntary donations has increased by 10% compared to the previous year, or if the proportion of voluntary donations within the total donations is 50%
- In addition, the blood banks will receive additional 3 GEL for each donation quarterly, if the proportion of voluntary donations within the total donations is 50%, and within these voluntary donations, 30% are regular voluntary donations



Progress Amendments to Licensing Requirements

The Safe Blood Working Group prepared the first draft of changes to licensing requirement.

Current Requirements	Amendments
Infrastructure and facilities for personal hygiene, cleaning and disinfection of the premises	Donor selection: All donors undergo a screening process to assess their suitability. Questionnaire must be designed to elicit information relevant to health and lifestyle of donor. On completion it is signed by the donor. List of conditions are in the standards and may be useful to include in the questionnaire. Hemoglobin concentration must be determined each time the donor attends to donate blood or blood components.
Safe medical waste segregation, collection, storage, disposal, recycling and/or destruction according to the rules established by legislation	Donation: - Minimum 12 square meters for donor chair and additionally, at least 6 square meters for extra chair - Scale for donation.
To ensure the proper regime of disinfection and sterilization and the relevant accounting according to the rules prescribed by legislation	Equipment: scale for donor, scale for blood components, tonometer, donor chair, refrigerated centrifuge, plasma extractor, domestic refrigerator, refrigerator for red cells (+2 +6, if produced), plasma freezer (-80 -16), Platelet incubator and shaker (+20 +24, if produced)



Progress Amendments to Licensing Requirements

Current Requirements	Amendments
Blood and blood components storage facilities and equipment according to the legislation	Infrastructure: registration and medical examination room with adequate furniture and facilities for registration and selection of donors, blood collection room, room for laboratory for blood group serology, room for laboratory for TTI testing (shall be air-conditioned), sterilization and washing room, refreshment room, waiting room, store and records room.
Uninterrupted water and electricity supply	Testing-typing: minimum ELISA for TTI testing and minimum Test Tube Method for blood group typing
Maintenance of medical and statistical documentation according to the legislation	Blood component preparation: refrigerated centrifuge, equipment for apheresis (for plasma centers)
Respective specialty certificates/qualifications of medical personnel as required by the legislation	Blood storage: storage areas must provide for appropriately secure and segregated storage of different types of blood and blood components including quarantine and released materials. Storage areas should be maintained within appropriate temperature limits. Frozen, refrigerated or heated storage conditions should have a temperature recording and alarm. Quarantined status areas should be marked clearly.
	Participation in the State External Quality Control System




Progress Activities

- All licensed blood banks and majority of hospitals performing blood transfusion are involved in the Donors' Unified Electronic Database enabling vein-to vein traceability of TTI infections
- Development of the national guidelines for blood donor TTI testing is in the process, including algorithms of blood donor screening as well as confirmatory testing, donor re-entry algorithms, product disposition, donor notification and management
- One laboratory specialist was contracted for providing external quality control and selection process for recruitment of transfusion medicine specialist is in progress
- Negotiation with Global Fund on procurement (donation) of two NAT equipment has been started



Progress Linkage to Diagnostic Services

- Confirmatory testing of HCV screening positive cases was introduced within the Hepatitis
 C Management Program that eliminated financial barrier to in depth diagnostic services
- Since 2018, blood banks transfer hepatitis C screening-positive samples to the Center for Cor-Ag confirmatory testing
- During January-February 2018, a total of 14608 donations were screened, 106 screening positive samples were sent to the Lugar Center, 55 confirmed results have been found





Progress KAP Survey

Voluntary
Unpaid Repeat
Donation

Main Barriers

Social problems, unclear procedures from donor registration to blood realization, donation stress

Lack or depreciation of important information, patients bear expensed for blood taken from voluntary donors

Remote location of blood banks, absence of advertisements and motivational stories in movies, TV series

Lack of actuality of the issue, believe that blood should not be sold at any cost, willingness popular faces deliver messages, rumors and myths

Mistrust to blood banks, lack of stimulation of unpaid voluntary repeat donations



Progress Communication Campaign

Activities

- Supporting events related to the World blood Donor Day were held within the week
- 4 blood drive days were held in corporative organizations (blood was taken from 230 persons)
- Popularization materials with campaign logotype and slogan were prepared and distributed among general and donor population
- A seminar for media representatives was held at NCDC
- Short motivational text messages were sent to 29,367 potential, eligible donors

Multimedia Campaign

- Live television broadcasting of voluntary blood donation was organized to raise public awareness
- 10 television broadcasts were dedicated to the popularization of voluntary repeat donations
- 24 infographics were prepared and posted on Facebook page I am a Blood Donor
- Number of creative and awareness raising posts were prepared and posted on Facebook page
 - 13 educational and informational articles were prepared and published in print and social media



Progress EU Technical Assistance Tools



In December 2017, NCDC submitted Strategic TAIEX and Twinning Projects to European Commission aiming at approximation of national blood safety regulations to EU directives.

DIRECTIVE 2002/98/EC OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 27 January 2003 setting standards of quality and safety for the collection, testing, processing, storage and distribution of human blood and blood components and amending Directive 2001/83/EC

COMMISSION DIRECTIVE 2004/33/EC of 22 March 2004 implementing Directive 2002/98/EC of the European Parliament and of the Council as regards certain technical requirements for blood and blood components

COMMISSION DIRECTIVE 2005/61/EC of 30 September 2005 implementing Directive 2002/98/EC of the European Parliament and of the Council as regards traceability requirements and notification of serious adverse reactions and events


COMMISSION DIRECTIVE 2005/62/EC of 30 September 2005 implementing Directive 2002/98/EC of the European Parliament and of the Council as regards Community standards and specifications relating to a quality system for blood establishments

- ★ TAIEX project was approved
- ★ Twinning project was included into the priorities short list





Challenges





Next Steps

Still need to be done

Finalize national guidelines for TTI testing

Finalize building of external quality control system and involve all blood banks

Finalize amendments to licensing requirements and submit for endorsement

Prepare for expert mission within TAIEX tool

Implement necessary activities to introduce NAT piloting in 2019 (including NAT apparatus and trainings)

Increase voluntary unpaid repeat donations

Conduct active communication campaign to raise population awareness on voluntary regular blood donations



Thank you for your attention!

Stay connected with us E-mail: e.kipiani@ncdc.ge Web site: donori.ncdc.ge



HCV treatment decentralization

Harm Reduction group

Maia Butsashvili

Background

- PWID are a major source of the burden of HCV in the country as documented by the national HCV sero-survey conducted in 2015
- Recent modeling indicates that priority treatment of PWID is needed to reach the elimination goals

Problems related with HCV infected PWID medical care

- Double-stigma in society
- Possible discrimination in medical institutions
- Unwillingness of medical infrastructure to meet the demands of this patient group
- Solution user friendly, non-judgmental and unbiased medical setting

Integrating HCV care with HR services

Integrated services

- Can mitigate stigma
- Encourage HCV treatment access by creating non-judgmental environment at the familiar institution (HR center)
- Based on trusting relationships and supports

The effectiveness of this strategy can be assessed to determine if this model can be expanded nationally.

Milestones of HCV treatment integration initiative with HR services

April, 2016

 Meeting on integration of HCV treatment with harm reduction services – during EASL meeting, Barcelona

June, 2016

- Meeting of ECHO and CDC team with civil society representatives and NCDC at HRU/NEOLAB office to discuss HCV treatment integration with HR services
- TAG recommendations

December 2017

First meeting on HCV treatment decentralization

Working group

Decentralization Effort

Objective: Decentralization of HCV treatment and careservices.

Oversight & Coordination Group

Objective: Coordination and management of the decentralization effort. Final decisions made by this group and approval sought from appropriate institutions

Primary Health Care/Hospital Group

Harm Reduction Group Objective: Development of a practical model of care to be implemented at designated health care sites. Presentation of plan to oversight group.

Challenges to Address

- Simplified Diagnostics and Treatment guidelines/algorithms
- Ensure Laboratory and Diagnostics integration
- Develop and Disseminate SOPs
- Develop Curriculum for Harm Reduction Centers
- Initial and Continued (CME) Training for Harm Reduction staff

Challenges to Address (cont.)

- Use of ECHO platform to facilitate training when applicable
- Develop necessary information systems/modules to support decentralization projects into the national Elim-C database
- Develop reasonable minimum criteria required for facilities to provide services to HCV patients
- Incorporate lessons, findings from harm reduction center related pilot projects into practical model/action plan.

Roles and Responsibilities:

- Meet as a group as recommended by the Oversight committee and the workgroup Chair
- Adhere to timeline of task completion outlined by the oversight committee and workgroup chair
- Develop an action plan for decentralization at Harm Reduction
 Centers that addresses the challenges below
- Adhere to expectations of meetings, communication, documents/products set by the oversight workgroup (meeting minutes, action plans, white papers, proposals, etc.)

Concerns related to integrating HCV treatment with HR services

- Unwillingness of HR center personnel
- Low trust among patients and preference to be treated at specialized clinics
- Quality of treatment and poorer outcome

Study to evaluate barriers of integrated care Funded by CDC/TEPHINET

- KAP survey of OST/NSP personnel on HCV care integration to evaluate readiness of health personnel to provide HCV care at OST/NSP sites.
- Qualitative evaluation of HCV treatment integration to understand barriers and challenges of HCV integrated care
- Patient satisfaction survey
- Chart review form for patients enrolled in integrated care services

Patient satisfaction survey

Preliminary results

Question	Answer	%
Is the location of medical facility convenient for you?	Yes	93.3
How satisfactory are the conditions (building, waiting space, sanitary conditions) at the medical facility?	Good	97.9
Are you satisfied with the attitude towards you from your doctor?	Yes	98
Are you satisfied with the attitude towards you from the nurses?	Yes	98

Patient satisfaction survey

Preliminary results

Have you received comprehensive information about hepatitis C treatment and side effects?	Yes	98%
Do you think that your confidentiality is protected during HCV treatment?	Yes	96.2%
Do you think that it's convenient to receive HCV treatment and methadone substitution therapy at the same clinic?	Yes	98%
In general, how would you assess HCV treatment services at this facility?	Very Good	93.5%
Will you recommend your family member/relative/friend (if interested) to be enrolled in National HCV Elimination Program?	Yes	100%

Provider attitude to HCV treatment integrated services (IS)

Preliminary data (N=51)

Question	Answer	%
Do you believe HCV treatment should be integrated with HR services	Yes	98
Do you believe your institution has enough administrative resources for IS?	Yes	82.8
Do you believe your institution has appropriate infrastructure for IS?	Yes	68.8
Willing to have IS at their institution	Yes	98


Diagnostics

Necessary diagnostic tests

- Screening
 - (HCV antibody test)
- Confirmatory testing
 - Core Antigen, RNA PCR (Quantitative, Qualitative), GeneExpert
 - Centralized location of confirmatory testing (Lugar Center)
- > Pre-treatment diagnostics
 - Simplify requirements
 - Minimize cost



Simplified Algorithm of pre-treatment diagnostics



Simplified Treatment Monitoring Algorithm

Simplified treatment monitoring procedures				
Measurements	Treatment Duration (weeks)			After treatment completion (weeks)
	4	8	12	12 or 24
Clinical assessment	X	Х	X	X
HCV RNA quantitative				X
Complete blood count	X*	X*	X*	
ALT	X	X	X	

^{*} only for patients receiving Ribavirin containing regimens

- Develop reasonable minimum criteria required for facilities to provide services to HCV patients
- ➤ Incorporate lessons, findings from pilot projects into model

